

Teva Pharmaceuticals (Pty) Ltd.

Astor 10/20/40/80 (film-coated tablets)

Each tablet contains 10/20/40/80 mg Atorvastatin as Atorvastatin calcium

PROFESSIONAL INFORMATION

SCHEDULING STATUS: S4

PROPRIETARY NAMES (AND DOSAGE FORM):

ASTOR 10 (film-coated tablets)

ASTOR 20 (film-coated tablets)

ASTOR 40 (film-coated tablets)

ASTOR 80 (film-coated tablets)

COMPOSITION:

ASTOR 10: Each film-coated tablet contains atorvastatin calcium equivalent to 10 mg atorvastatin.

ASTOR 20: Each film-coated tablet contains atorvastatin calcium equivalent to 20 mg atorvastatin.

ASTOR 40: Each film-coated tablet contains atorvastatin calcium equivalent to 40 mg atorvastatin.

ASTOR 80: Each film-coated tablet contains atorvastatin calcium equivalent to 80 mg atorvastatin.

ASTOR 10, 20, 40 & 80 film-coated tablets also contain the following inactive ingredients: activated attapulgate, Microcelac 100 (consisting of lactose monohydrate and microcrystalline cellulose), microcrystalline cellulose, pregelatinised starch, hydroxypropyl cellulose, magnesium stearate, silica colloidal anhydrous, and Opadry White (consisting of hypromellose, lactose monohydrate, polyethylene glycol and titanium dioxide).

PHARMACOLOGICAL CLASSIFICATION:

A 7.5 Serum-cholesterol reducers.

Teva Pharmaceuticals (Pty) Ltd.

Astor 10/20/40/80 (film-coated tablets)

Each tablet contains 10/20/40/80 mg Atorvastatin as Atorvastatin calcium

PHARMACOLOGICAL ACTION:

Atorvastatin is a fully synthetic cholesterol-lowering agent. It is a competitive inhibitor of 3-hydroxy-3-methylglutaryl coenzyme A (HMG-CoA) reductase, the enzyme that catalyses the conversion of HMG-CoA to mevalonate, an early and rate-limiting step in the biosynthesis of cholesterol. As a result, in patients with hyperlipidaemias, including hypercholesterolaemias and combined (mixed) hyperlipidaemia (type IIa or IIb hyperlipoproteinaemias), hypertriglyceridaemia (type IV), and dysbetalipoproteinaemia (type III), atorvastatin reduces total plasma cholesterol, low-density lipoprotein (LDL-C)- and very low density lipoprotein (VLDL-C)-cholesterol concentrations, triglycerides and apolipoprotein B. In addition, atorvastatin increases high-density lipoprotein (HDL-C)-cholesterol. Atorvastatin exerts its main effect in the liver.

Pharmacokinetics:

Absorption:

Atorvastatin is administered in an active (open) form without the need for hydrolysis. It is rapidly absorbed from the gastrointestinal tract, with maximum concentration being achieved in 1 to 2 hours. It has a low absolute bioavailability of the parent compound of about 12 %, due to presystemic clearance in the gastrointestinal mucosa and/or first-pass metabolism in the liver, its primary site of action.

LDL-C reduction is similar whether atorvastatin is given with or without food, despite a decreased rate and extent of absorption when given with food.

LDL-C reduction is the same regardless of the time of administration, despite lower atorvastatin plasma concentrations following evening administration compared to morning administration.

Distribution:

Atorvastatin is 98 % or more bound to plasma proteins.

Metabolism:

Atorvastatin is metabolised by the cytochrome P450 isoenzyme CYP3A4 to a number of active metabolites (ortho- and parahydroxylated derivatives and various beta-oxidation products). The

Teva Pharmaceuticals (Pty) Ltd.

Astor 10/20/40/80 (film-coated tablets)

Each tablet contains 10/20/40/80 mg Atorvastatin as Atorvastatin calcium

ortho- and parahydroxylated metabolites are pharmacologically active and their in vitro inhibition of HMG-CoA reductase activity is equivalent to the parent compound.

Elimination:

The mean plasma elimination half-life of atorvastatin is about 14 hours, although the half-life of inhibitory activity for HMG-CoA reductase is about 20 to 30 hours due to the contribution of the active metabolites. Atorvastatin is excreted as metabolites, primarily in the bile. Less than 2 % is excreted renally.

Special populations:

Elderly:

Lipid lowering effects in the elderly are comparable to those in young adults at equal doses, despite higher plasma concentrations of atorvastatin in healthy elderly subjects.

Paediatrics:

Pharmacokinetic data in the paediatric population are not available.

Hepatic impairment:

Plasma concentrations of atorvastatin are markedly increased (approximately 16-fold in C_{max} and 11-fold in AUC) in patients with chronic alcoholic liver disease (Child-Pugh B); see **CONTRA-**

INDICATIONS.

Renal impairment:

Renal impairment has no influence on plasma concentrations; therefore dose adjustment is not needed; see **WARNINGS.**

INDICATIONS:

ASTOR is indicated, in combination with diet, to decrease elevated total cholesterol, LDL-cholesterol, apolipoprotein-B and triglyceride levels in patients with:

Teva Pharmaceuticals (Pty) Ltd.

Astor 10/20/40/80 (film-coated tablets)

Each tablet contains 10/20/40/80 mg Atorvastatin as Atorvastatin calcium

- primary hypercholesterolaemia,
- heterozygous familial hypercholesterolaemia, and
- mixed dyslipidaemia.

ASTOR is also indicated to reduce total-C and LDL-C in patients with homozygous familial hypercholesterolaemia, as an adjunct to other lipid-lowering treatments (e.g. LDL apheresis), of if such treatments are not available.

Therapy with lipid-lowering agents should be a component of multiple-risk-factor intervention in individuals at increased risk of atherosclerotic vascular disease due to hypercholesterolaemia. Lipid-altering agents should be used in addition to a diet restricted in saturated fat and cholesterol only when the response to diet and other non-pharmacological measures has been inadequate.

Secondary causes for hypercholesterolaemia (e.g. poorly controlled diabetes mellitus, nephrotic syndrome, hypothyroidism, obstructive liver disease, dysproteinaemias, alcoholism and therapy with other medicines) should be excluded prior to initiating therapy with ASTOR, and a lipid profile performed to measure total-C, LDL-C, HDL-C and triglycerides.

CONTRA-INDICATIONS:

- Hypersensitivity to atorvastatin, other HMG-CoA reductase inhibitors or any component of ASTOR.
- Active hepatic disease (the condition may be exacerbated) or unexplained persistently raised serum-
- aminotransferase concentrations (exceeding 3 times the upper limit of normal).
- Pregnancy and lactation (see **PREGNANCY AND LACTATION**).
- Patients with Child-Pugh B and C liver impairment.
- Concomitant use with rifampicin, diltiazem and grapefruit juice (see **INTERACTIONS**).

WARNINGS:

Effects on the liver:

Serum transaminase:

Teva Pharmaceuticals (Pty) Ltd.

Astor 10/20/40/80 (film-coated tablets)

Each tablet contains 10/20/40/80 mg Atorvastatin as Atorvastatin calcium

Serum transaminase values may be increased, usually to less than 3 times the upper limit of normal, in slightly less than 1 to 2 % of patients receiving HMG-CoA reductase inhibitors for at least 1 year. Marked increases to more than 3 times the upper limit of normal have occurred.

Liver function tests, including serum transaminase:

Determinations are recommended prior to initiation of therapy, following each dosage increase, and subsequently when clinically indicated. ASTOR should be discontinued if the rise in transaminase levels is persistent and/or increases to three times the upper limit of normal (ULN) or more.

ASTOR should be used with caution in patients who consume substantial amounts of alcohol and/or who have a history of liver disease.

Renal impairment:

Renal impairment has no influence on plasma concentrations; therefore dose adjustment is not needed. (See also rhabdomyolysis under Skeletal muscle below).

Skeletal muscle:

ASTOR may cause myopathy and rhabdomyolysis, especially at higher doses, and it should be used with caution in patients at risk of rhabdomyolysis, and particularly in patients taking medicines, such as cytochrome P450 inhibitors (see **INTERACTIONS**), that increase plasma concentrations of the HMG-CoA reductase inhibitor, ASTOR.

Rhabdomyolysis with or without renal impairment has been reported with the use of HMG-CoA reductase inhibitors such as ASTOR. A history of renal impairment may be a risk factor for the development of rhabdomyolysis. Such patients merit closer monitoring for the skeletal muscle adverse events.

General measures to reduce the risk of myopathy:

Patients starting treatment with ASTOR should be advised of the risk of myopathy and should promptly report unexplained muscle pain, tenderness, or weakness, especially if accompanied by malaise or fever. A creatine kinase (CK) level above 10 times the Upper Limit of Normal (ULN) in a

Teva Pharmaceuticals (Pty) Ltd.

Astor 10/20/40/80 (film-coated tablets)

Each tablet contains 10/20/40/80 mg Atorvastatin as Atorvastatin calcium

patient, with unexplained symptoms, indicate myopathy. ASTOR should be discontinued if creatine phosphokinase increases significantly or if myopathy is diagnosed.

Measures to reduce the risk of myopathy caused by medicine interactions:

The benefits of using ASTOR concomitantly with immunosuppressants, fibrates or lipid-lowering doses of niacin should be carefully considered. Concomitant administration with ciclosporin, itraconazole, ketoconazole, erythromycin, clarithromycin, HIV-protease inhibitors and nefazodone is not recommended. In patients receiving ciclosporin, ASTOR should be temporarily discontinued.

Haemorrhagic stroke:

Patients without coronary heart disease who had a stroke or transient ischaemic attack (TIA) within the preceding months who were initiated on atorvastatin 80 mg revealed a higher incidence of haemorrhagic stroke compared to placebo. Patients with haemorrhagic stroke on entry appeared to be at risk for recurrent haemorrhagic stroke.

Paediatric patients:

Use in paediatric patients is not recommended, as safety and efficacy have not been established.

INTERACTIONS:

Inhibitors of cytochrome P450:

Atorvastatin is metabolised by cytochrome P450 3A4. Concomitant administration of ASTOR with inhibitors of cytochrome P450 3A4 can lead to an increase in plasma concentrations of atorvastatin. Medicines that inhibit cytochrome P450 isoenzyme CYP3A4 include: ciclosporin, itraconazole, ketoconazole, erythromycin/clarithromycin (see **Macrolides** below), HIV-protease inhibitors, amiodarone, verapamil and nefazodone. There is a similar interaction with grapefruit juice (contraindicated); see **Grapefruit juice** below.

Macrolides:

Teva Pharmaceuticals (Pty) Ltd.

Astor 10/20/40/80 (film-coated tablets)

Each tablet contains 10/20/40/80 mg Atorvastatin as Atorvastatin calcium

Erythromycin/clarithromycin: In healthy individuals plasma concentrations of atorvastatin increased approximately 40 % with co-administration of erythromycin, a known inhibitor of CYP 3A4.

Azithromycin: Co-administration of atorvastatin 10 mg and azithromycin (500 mg once daily) did not alter the plasma concentrations of atorvastatin.

Grapefruit juice:

Co-administration of grapefruit juice and atorvastatin may increase the concentration of atorvastatin, as in ASTOR, by 2,5 to 3,3 fold. Therefore the combination should be avoided (see **CONTRA-INDICATIONS**).

Inducers of cytochrome P450 3A4:

Concomitant administration of atorvastatin with inducers of cytochrome P450 3A4, such as efavirenz and rifampicin can lead to variable reductions in plasma concentrations of atorvastatin. Due to the dual mechanism of rifampicin, simultaneous co-administration of ASTOR with rifampicin is not recommended, as delayed administration of atorvastatin after administration of rifampicin has been associated with a significant reduction in atorvastatin plasma concentrations (see **CONTRA-INDICATIONS**).

Cimetidine:

Co-administration of atorvastatin with cimetidine does not alter plasma concentration and LDL reduction.

Diltiazem HCl:

Co-administration of atorvastatin with diltiazem was associated with a 51 % increase in the AUC of atorvastatin. Therefore the combination should be avoided (see **CONTRA-INDICATIONS**).

Fucidic acid:

Teva Pharmaceuticals (Pty) Ltd.

Astor 10/20/40/80 (film-coated tablets)

Each tablet contains 10/20/40/80 mg Atorvastatin as Atorvastatin calcium

Severe muscle problems such as rhabdomyolysis have been reported with the concomitant use of fucidic acid and atorvastatin. Patients on fucidic acid and ASTOR should be closely monitored and temporary suspension of ASTOR may be appropriate.

Transporter Inhibitors:

Inhibitors of the OATP1B1 (organic anion-transporting polypeptide-1B1) transport system, such as ciclosporin, can increase the bioavailability of atorvastatin. Concomitant administration of atorvastatin 10 mg and ciclosporin 5,2 mg/kg per day resulted in a 7,7 fold increase in exposure to atorvastatin.

Warfarin:

A possible increase in the anticoagulant effect of warfarin may occur. Patients taking warfarin should have their INR determined before starting ASTOR therapy. The INR should be monitored frequently enough in the early stages of therapy until stabilised. Once a stable INR has been documented, INR can be monitored at the intervals usually recommended for patients on warfarin. When there is a dose adjustment of ASTOR, this procedure should be repeated.

Digoxin:

Concurrent use may cause an elevation in serum digoxin concentrations by approximately 20 %.

Bile acid sequestrants:

ASTOR should be taken 1 hour before or 4 hours after cholestyramine. Concurrent use may decrease the bioavailability of ASTOR.

Azole antifungals, ciclosporin, gemfibrozil, other fibrates, immunosuppressants, macrolide antibiotics or niacin:

Concurrent use with ASTOR may be associated with an increased risk of myopathy, myositis, rhabdomyolysis and acute renal failure (see **WARNINGS**).

Antacids:

Teva Pharmaceuticals (Pty) Ltd.

Astor 10/20/40/80 (film-coated tablets)

Each tablet contains 10/20/40/80 mg Atorvastatin as Atorvastatin calcium

Concurrent use may decrease plasma concentrations of atorvastatin by approximately 35 %. LDL-C reduction is however not altered.

Oral contraceptives:

Concurrent use with atorvastatin may increase the AUC value for norethindrone and ethinyl oestradiol by approximately 30 % and 20 % respectively.

PREGNANCY AND LACTATION:

Safety and efficacy in pregnancy and lactation have not been established. The use of ASTOR during pregnancy and lactation, or in women who plan to become pregnant, is contra-indicated.

Women of child-bearing potential should use appropriate contraceptive measures. In the event of planning a pregnancy, an interval of one month should be allowed from stopping ASTOR treatment.

DOSAGE AND DIRECTIONS FOR USE:

The patient must follow a cholesterol-lowering diet before initiation of, and while on ASTOR therapy.

ASTOR can be taken at any time of the day with meals or on an empty stomach.

ASTOR should not be taken with grapefruit juice.

Hypercholesterolaemia:

Heterozygous familial hypercholesterolaemia and mixed dyslipidaemia:

The usual starting dose is 10 mg of ASTOR daily. The dose may be adjusted at intervals of 4 weeks up to a maximum of 80 mg daily.

Children:

Treatment experience in the paediatric population is limited.

Homozygous familial hypercholesterolaemia:

10 to 80 mg ASTOR once per day. ASTOR should be used in these patients as an adjunct to other lipid-lowering treatments such as LDL apheresis, or if such treatments are unavailable.

Teva Pharmaceuticals (Pty) Ltd.

Astor 10/20/40/80 (film-coated tablets)

Each tablet contains 10/20/40/80 mg Atorvastatin as Atorvastatin calcium

Prevention of cardiovascular complications:

The usual dose is 10 mg ASTOR once per day.

Dosage in patients with renal insufficiency:

Dosage adjustment in patients with renal dysfunction is not necessary because renal disease does not affect the plasma concentrations nor LDL-C reduction (however, see **WARNINGS**).

Dosage in patients with hepatic impairment:

In patients with moderate to severe hepatic dysfunction, the therapeutic response to ASTOR is unaffected, but serum levels of the medicine are significantly increased. In patients with chronic alcoholic liver disease, plasma concentrations of atorvastatin are markedly increased. Both C_{max} and AUC are 4-fold greater in patients with Child-Pugh A disease. C_{max} and AUC are approximately 16-fold and 11-fold increased, respectively, in patients with Child-Pugh B disease. Therefore, caution with dosing should be exercised in patients who take substantial quantities of alcohol and/or have a history of liver disease (see **CONTRA-INDICATIONS** and **WARNINGS**).

SIDE-EFFECTS AND SPECIAL PRECAUTIONS:

Side-Effects:

Blood and lymphatic system disorders:

Less frequent: Thrombocytopenia, anaemia, neutropenia.

Immune system disorders:

Frequent: Hypersensitivity reactions, including anaphylaxis and angioedema.

Metabolism and nutrition disorders:

Less frequent: Hypoglycaemia, hyperglycaemia, weight gain, anorexia.

Psychiatric disorders:

Frequent: Insomnia.

Teva Pharmaceuticals (Pty) Ltd.

Astor 10/20/40/80 (film-coated tablets)

Each tablet contains 10/20/40/80 mg Atorvastatin as Atorvastatin calcium

Nervous system disorders:

Frequent: Dizziness, headache, paraesthesia, hypoaesthesia.

Less frequent: Peripheral neuropathy, cognitive impairment such as memory loss, forgetfulness, amnesia, memory impairment and confusion.

Frequency unknown: Myasthenia gravis.

Eye disorders:

Less frequent: Blurred vision, visual disturbance.

Frequency unknown: Ocular myasthenia.

Ear and labyrinth disorders:

Less frequent: Tinnitus, hearing loss.

Vascular disorders:

Less frequent: Peripheral oedema.

Respiratory, thoracic and mediastinal disorders:

Frequency unknown: Sinusitis, pharyngitis.

Gastrointestinal disorders:

Frequent: Constipation, diarrhoea, flatulence, heartburn, abdominal pain and cramps, nausea, dyspepsia.

Less frequent: Dysgeusia (taste disturbances), vomiting, pancreatitis, anorexia.

Hepatobiliary disorders:

Less frequent: Hepatitis, cholestatic jaundice, hepatic failure.

Skin and subcutaneous tissue disorders:

Teva Pharmaceuticals (Pty) Ltd.

Astor 10/20/40/80 (film-coated tablets)

Each tablet contains 10/20/40/80 mg Atorvastatin as Atorvastatin calcium

Frequent: Skin rash, pruritus.

Less frequent: Alopecia, Stevens-Johnson syndrome, toxic epidermal necrolysis, erythema multiforme, bullous rashes, urticaria.

Musculoskeletal, connective tissue and bone disorders:

Frequent: Myalgia, arthralgia, back pain.

Less frequent: Myopathy, characterised by myalgia and muscle weakness, and associated with increased creatine phosphokinase concentrations, rhabdomyolysis with acute renal failure, myositis, muscle cramps, tendon rupture.

Reproductive system and breast disorders:

Less frequent: Impotence (decreased sexual ability), gynaecomastia.

General disorders and administration site conditions:

Frequent: Fatigue, asthenia, chest pain, infection.

Less frequent: Malaise.

Laboratory test findings:

Marked and persistent increases of serum transaminases and elevated alkaline phosphatase and gamma-glutamyl transpeptidase have been reported. Liver function test abnormalities have generally been mild and transient. Increases in serum creatinine kinase (CK) levels, derived from skeletal muscle, have been reported (see **Special Precautions**).

Special precautions:

ASTOR should be used with caution in:

- Patients who consume substantial amounts of alcohol and/or who have a history of liver disease.
- Liver enzyme tests should be performed in patients before initiating ASTOR therapy and as clinically indicated thereafter. If serious liver injury with clinical symptoms and/or

Teva Pharmaceuticals (Pty) Ltd.

Astor 10/20/40/80 (film-coated tablets)

Each tablet contains 10/20/40/80 mg Atorvastatin as Atorvastatin calcium

hyperbilirubinaemia or jaundice occurs during treatment, therapy should be interrupted. If an alternate aetiology is not found, ASTOR should not be restarted.

- Patients who may be predisposed to developing renal failure secondary to rhabdomyolysis such as those with severe acute infection, hypotension, severe metabolic, endocrine or electrolyte disorders, uncontrolled seizures, major surgery or trauma. There is an increased risk of developing renal failure if rhabdomyolysis occurs. The onset of rhabdomyolysis may occur weeks to months after initiation of treatment (see **WARNINGS**).
- Patients who have severe renal impairment.
- Risk of myasthenia gravis and ocular myasthenia.

Creatine kinase (CK) concentrations:

Mild transient increases are common and may not be medication related. Medication related marked increases, with myositis and possible renal failure occur in about 0,5 to 1 % of patients, although the incidence may be higher in organ transplant patients treated concurrently with immunosuppressants or gemfibrozil.

Determination of serum creatine kinase is recommended if the patient develops muscle tenderness during therapy or during concurrent therapy with niacin or immunosuppressive medications. A level of 10 times higher than the upper limit of normal in a patient with unexplained muscle symptoms indicates myopathy.

Organ transplant with immunosuppressive therapy:

Increased risk of rhabdomyolysis and renal failure.

Lactose intolerance:

ASTOR contains lactose. Patients who are lactose intolerant or have rare hereditary problems of galactose intolerance, Lapp lactase deficiency or glucose-galactose malabsorption should not take ASTOR.

Ability to drive or operate machinery:

Teva Pharmaceuticals (Pty) Ltd.

Astor 10/20/40/80 (film-coated tablets)

Each tablet contains 10/20/40/80 mg Atorvastatin as Atorvastatin calcium

ASTOR may cause blurred vision, dizziness and confusion.

Patients should therefore not operate hazardous machinery, including motor vehicles, until they are reasonably certain that ASTOR does not adversely affect them.

KNOWN SYMPTOMS OF OVERDOSAGE AND PARTICULARS OF ITS TREATMENT:

(See **Side-Effects and Special Precautions**).

General measures should be adopted and liver function should be monitored.

Treatment is symptomatic and supportive. Haemodialysis is not expected to significantly increase atorvastatin (as in ASTOR) clearance, due to extensive plasma protein binding.

IDENTIFICATION:

ASTOR 10: White oval biconvex film-coated tablets, approximately 10,1 x 5,6 mm and 3,7 mm in thickness.

ASTOR 20: White oval biconvex film-coated tablets with a non-functional breakline on one side, approximately 12,7 x 6,7 mm and 4,6 mm in thickness.

ASTOR 40: White oblong biconvex film-coated tablets, approximately 19,4 x 7,8 mm and 4,7 mm in thickness.

ASTOR 80: White oblong biconvex film-coated tablets with a non-functional breakline on one side, approximately 22,8 x 10,9 mm and 5,7 mm in thickness.

PRESENTATION:

Packs of 28 (4 blisters of 7 tablets) or 30 (3 blisters of 10 tablets) film-coated tablets packed in Aluminium/Aluminium blister strips in a cardboard carton.

Not all pack sizes are necessarily marketed.

STORAGE INSTRUCTIONS:

Store in the original packaging (in the carton) at or below 25 °C.

KEEP OUT OF REACH OF CHILDREN.

Teva Pharmaceuticals (Pty) Ltd.

Astor 10/20/40/80 (film-coated tablets)

Each tablet contains 10/20/40/80 mg Atorvastatin as Atorvastatin calcium

REGISTRATION NUMBERS:

ASTOR 10: 44/7.5/1054

ASTOR 20: 44/7.5/1055

ASTOR 40: 44/7.5/1056

ASTOR 80: 44/7.5/1057

NAME AND BUSINESS ADDRESS OF THE HOLDER OF THE CERTIFICATES OF

REGISTRATION:

Teva Pharmaceuticals (Pty) Ltd,

Maxwell Office Park

Magwa Crescent West

Waterfall City

Midrand

Gauteng

2090

South Africa

DATE OF PUBLICATION OF THIS PACKAGE INSERT:

30 September 2025