

## PROFESSIONAL INFORMATION

### SCHEDULING STATUS

S4

#### 1. NAME OF THE MEDICINE

Lipocomb 10 mg/10 mg hard gelatin capsules.

Lipocomb 20 mg/10 mg hard gelatin capsules.

#### 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Lipocomb 10 mg/10 mg hard gelatin capsules.

Each capsule contains 10 mg rosuvastatin (as zinc) and 10 mg of ezetimibe.

Lipocomb 20 mg/10 mg hard gelatin capsules.

Each capsule contains 20 mg rosuvastatin (as zinc) and 10 mg of ezetimibe.

Excipient with known effect: mannitol.

Lipocomb 10 mg/10 mg contains 30,3 mg mannitol.

Lipocomb 20 mg/10 mg contains 30,3 mg mannitol.

For the full list of excipients, see section 6.1.

#### 3. PHARMACEUTICAL FORM

Hard Gelatin Capsules.

Lipocomb 10 mg/10 mg hard gelatin capsules: Unmarked self-closing Coni Snap type, size 0, hard gelatin capsule with yellow coloured cap and yellow coloured body filled with two tablets. The length of the capsule is about 21,7 mm ( $\pm$  0,5 mm).

Lipocomb 20 mg/10 mg hard gelatin capsules: Unmarked self-closing Coni Snap type, size 0, hard gelatin capsule with caramel coloured cap and yellow coloured body filled with two tablets. The length of the capsule is about 21,7 mm ( $\pm$  0,5 mm).

#### 4. CLINICAL PARTICULARS

##### 4.1 Therapeutic indications

Lipocomb is indicated as adjunct to diet for treatment of primary hypercholesterolemia as substitution therapy in adult patients adequately controlled with the individual substances given concurrently at the same dose level as in the fixed dose combination, but as separate products.

## **4.2 Posology and method of administration**

### **Posology**

Lipocomb is indicated in adult patients whose hypercholesterolemia is adequately controlled with separately administered monocomponent preparations of the same doses as the recommended combination.

The patient should be on an appropriate lipid-lowering diet and should continue on this diet during treatment with Lipocomb.

The recommended daily dose is one capsule daily.

Lipocomb is not suitable for initial therapy. Treatment initiation or dose adjustment if necessary should only be done with the monocomponents and after setting the appropriate doses the switch to the fixed dose combination of the appropriate strength is possible.

Lipocomb 10 mg/10 mg and 20 mg/10 mg hard gelatin capsules are not suitable for the treatment of patients requiring 40 mg dose of rosuvastatin.

### **Special populations**

#### **Elderly**

A start dose of 5 mg rosuvastatin is recommended in patients > 70 years (see section 4.4).

The combination is not suitable for initial therapy. Treatment initiation or dose adjustment if necessary, should only be done with the monocomponents and after setting the appropriate doses the switch to the fixed dose combination of the appropriate strength is possible.

#### **Renal impairment**

No dose adjustment is necessary in patients with mild to moderate renal impairment.

The recommended start dose is rosuvastatin 5 mg in patients with moderate renal impairment (creatinine clearance < 60 ml/min). The fixed dose combination is not suitable for initial therapy. Monocomponent preparations should be used to start the treatment or to modify the dose.

The use of Lipocomb in patients with severe renal impairment is contraindicated for all doses (see sections 4.3 and 5.2).

#### **Hepatic impairment**

No dose adjustment is required in patients with mild hepatic impairment (Child Pugh score 5 to 6). Treatment with Lipocomb is not recommended in patients with moderate (Child Pugh score 7 to 9) or severe (Child Pugh score > 9) liver dysfunction (see sections 4.4 and 5.2). Lipocomb is contraindicated in patients with active liver disease (see section 4.3).

### **Race**

Increased systemic exposure of rosuvastatin has been seen in Asian subjects (see sections 4.4 and 5.2). The recommended start dose is rosuvastatin 5 mg for patients of Asian ancestry. The fixed dose combination is not suitable for initial therapy. Monocomponent preparations should be used to start the treatment or to modify the dose.

### **Genetic polymorphisms**

Specific types of genetic polymorphisms are known that can lead to increased rosuvastatin exposure (see section 5.2). For patients who are known to have such specific types of polymorphisms, a lower daily dose of Lipocomb is recommended.

### **Dosage in patients with pre-disposing factors to myopathy**

The recommended start dose is rosuvastatin 5 mg in patients with predisposing factors to myopathy (see section 4.4). The fixed dose combination is not suitable for initial therapy. Monocomponent preparations should be used to start the treatment or to modify the dose.

### **Concomitant therapy**

Rosuvastatin is a substrate of various transporter proteins (e.g. OATP1B1 and BCRP). The risk of myopathy (including rhabdomyolysis) is increased when Lipocomb is administered concomitantly with certain medicines that may increase the plasma concentration of rosuvastatin due to interactions with these transporter proteins (certain protease inhibitors including combinations of ritonavir with atazanavir, lopinavir, and/or tipranavir; see sections 4.4 and 4.5). Whenever possible, alternative medications should be considered, and, if necessary, consider temporarily discontinuing Lipocomb therapy. In situations where co-administration of these medicines with Lipocomb is unavoidable, the dose of Lipocomb should not exceed 10 mg + 10 mg once daily.

### **Dosage in patients taking other medicines**

Lipocomb should be taken either  $\geq 2$  hours before or  $\geq 4$  hours after administration of a bile acid sequestrant.

#### **Co-administration with elbasvir or grazoprevir**

The dose of Lipocomb should not exceed 10 mg + 10 mg once daily.

#### **Co-administration with gemfibrozil**

Increased systemic exposure to rosuvastatin has been observed in patients taking concomitant rosuvastatin and gemfibrozil. If Lipocomb is taken in combination with gemfibrozil, the dose of Lipocomb should be limited to 10 mg + 10 mg once daily.

#### **Paediatric population**

The safety and efficacy of Lipocomb in children below the age of 18 years have not yet been established. Currently available data are described in sections 4.8, 5.1 and 5.2 but no recommendation on a posology can be made.

#### **Method of administration**

For oral use.

Lipocomb should be taken each day once at the same time of the day with or without food. The capsule should be swallowed whole with a drink of water.

#### **4.3 Contraindications**

Lipocomb is contraindicated:

- in patients with hypersensitivity to the active substances (rosuvastatin, ezetimibe) or to any of the excipients listed in section 6.1.
- in patients with active liver disease including unexplained, persistent elevations of serum transaminases and any serum transaminase elevation exceeding 3 times the upper limit of normal (ULN).
- during pregnancy and breast-feeding and in women of childbearing potential not using appropriate contraceptive measures.
- in patients with severe renal impairment (creatinine clearance  $< 30$  ml/min).
- moderate to severe hepatic impairment (Child Pugh score 7 or more).

- in patients with myopathy.
- in patients receiving concomitant ciclosporin. (see sections 4.4, 4.5 and 5.2).
- combination with fenofibrate in patients with gall bladder disease (see section 4.4).

#### **4.4 Special warnings and precautions for use**

##### **Skeletal Muscle Effects**

Effects on skeletal muscle e.g. myalgia, myopathy, and, rarely rhabdomyolysis have been reported in rosuvastatin-treated patients with all doses and in particular with doses > 20 mg.

In post-marketing experience with ezetimibe, cases of myopathy and rhabdomyolysis have been reported. However, rhabdomyolysis has been reported with ezetimibe monotherapy and with the addition of ezetimibe to other medicines known to be associated with increased risk of rhabdomyolysis. If myopathy is suspected based on muscle symptoms or is confirmed by a creatine kinase level, ezetimibe, any statin, and any of these medicines known to be associated with increased risk of rhabdomyolysis, that the patient is taking concomitantly should be immediately discontinued. All patients starting should be told to report promptly any unexplained muscle pain, tenderness or weakness (see section 4.8).

##### **Creatine Kinase Measurement**

Creatine kinase (CK) should not be measured following strenuous exercise or in the presence of a plausible alternative cause of CK increase, which may confound interpretation of the results.

If CK levels are significantly elevated at baseline (> 5 x ULN) a confirmatory test should be carried out within 5 - 7 days. If the repeat test confirms a baseline CK > 5 x ULN, treatment should not be started.

##### **Before treatment**

**Lipocomb**, as other HMG-CoA reductase inhibitors, should be prescribed with caution in patients with pre-disposing factors for myopathy/rhabdomyolysis. Such factors include:

- renal impairment
- hypothyroidism
- personal or family history of hereditary muscular disorders
- previous history of muscular toxicity with another HMG-CoA reductase inhibitor or fibrate
- alcohol abuse
- age > 70 years
- situations where an increase in plasma levels may occur (see section 5.2)
- concomitant use of fibrates.

In such patients the risk of treatment should be considered in relation to possible benefit and clinical monitoring is recommended. If CK levels are significantly elevated at baseline (> 5 x ULN) treatment should not be started.

#### **Whilst on treatment**

Patients should be asked to report inexplicable muscle pain, weakness or cramps immediately, particularly if associated with malaise or fever. CK levels should be measured in these patients. Therapy should be discontinued if CK levels are markedly elevated (> 5 x ULN) or if muscular symptoms are severe and cause daily discomfort (even if CK levels are > 5 x ULN). Routine monitoring of CK levels in asymptomatic patients is not warranted.

There have been reports of an immune-mediated necrotising myopathy (IMNM) during or after treatment with statins, including rosuvastatin. IMNM is clinically characterized by proximal muscle weakness and elevated serum creatine kinase, which persist despite discontinuation of statin treatment.

In clinical trials there was no evidence of increased skeletal muscle effects in the small number of patients dosed with rosuvastatin and concomitant therapy. However, an increase in the incidence of myositis and myopathy has been seen in patients receiving other HMG-CoA reductase inhibitors together with fibric acid derivatives including gemfibrozil, ciclosporin, nicotinic acid, azole antifungals, protease inhibitors and macrolide

antibiotics. Gemfibrozil increases the risk of myopathy when given concomitantly with some HMG-CoA reductase inhibitors. Therefore, the combination of Lipocomb and gemfibrozil is not recommended.

Lipocomb should not be used in any patient with an acute, serious condition suggestive of myopathy or predisposing to the development of renal failure secondary to rhabdomyolysis (e.g. sepsis, hypotension, major surgery, trauma, severe metabolic, endocrine and electrolyte disorders; or uncontrolled seizures).

### **Fusidic acid**

Lipocomb must not be co-administered with systemic formulations of fusidic acid or within 7 days of stopping fusidic acid treatment. In patients where the use of systemic fusidic acid is considered essential, statin treatment should be discontinued throughout the duration of fusidic acid treatment. There have been reports of rhabdomyolysis (including some fatalities) in patients receiving fusidic acid and statins in combination (see section 4.5). The patient should be advised to seek medical advice immediately if they experience any symptoms of muscle weakness, pain or tenderness.

Statin therapy may be re-introduced seven days after the last dose of fusidic acid.

In exceptional circumstances, where prolonged systemic fusidic acid is needed, e.g., for the treatment of severe infections, the need for co-administration of Lipocomb and fusidic acid should only be considered on a case by case basis and under close medical supervision.

### **Liver effects**

In patients receiving ezetimibe with a statin, consecutive transaminase elevations ( $\geq 3$  times the upper limit of normal [ULN]) have been observed.

It is recommended that liver functions tests be carried out 3 months following the initiation of rosuvastatin treatment. Rosuvastatin should be discontinued or the dose reduced if the level of serum transaminases is greater than 3 times the upper limit of normal.

In patients with secondary hypercholesterolaemia caused by hypothyroidism or nephrotic syndrome, the underlying disease should be treated prior to initiating therapy with Lipocomb.

Due to the unknown effects of the increased exposure to ezetimibe in patients with moderate or severe hepatic impairment, Lipocomb is not recommended (see section 5.2).

### **Renal effects**

Proteinuria, detected by dipstick testing and mostly tubular in origin, has been observed in patients treated with higher doses of rosuvastatin, in particular 40 mg, where it was transient or intermittent in most cases. Proteinuria has not been shown to be predictive of acute or progressive renal disease (see section 4.8).

### **Race**

Rosuvastatin pharmacokinetic studies show an increase in exposure in Asian subjects compared with Caucasians (see sections 4.2 and 5.2).

### **Protease inhibitors**

Increased systemic exposure to rosuvastatin has been observed in subjects receiving rosuvastatin concomitantly with various protease inhibitors in combination with ritonavir. Consideration should be given both to the benefit of lipid lowering by use of Lipocomb in HIV patients receiving protease inhibitors and the potential for increased rosuvastatin plasma concentrations when initiating and up titrating rosuvastatin in patients treated with protease inhibitors. The concomitant use with certain protease inhibitors is not recommended unless the dose of Lipocomb is adjusted (see sections 4.2 and 4.5).

### **Interstitial lung disease**

Exceptional cases of interstitial lung disease have been reported with some statins, especially with long term therapy (see section 4.8). Presenting features can include dyspnoea, non-productive cough and deterioration in general health (fatigue, weight loss and fever). If it is suspected a patient has developed interstitial lung disease, statin therapy should be discontinued.

### **Diabetes mellitus**

Increases in HbA1c and serum glucose levels have been observed in patients treated with rosuvastatin. An increased frequency of diabetes mellitus has been reported with rosuvastatin, as in Lipocomb, in patients with risk factors for diabetes mellitus

Patients at risk (fasting glucose 5,6 to 6,9 mmol/L, BMI > 30 kg/m<sup>2</sup>, raised triglycerides, hypertension) should be monitored both clinically and biochemically according to national guidelines.

### **Fibrates**

The safety and efficacy of ezetimibe administered with fibrates have not been established. If cholelithiasis is suspected in a patient receiving Lipocomb and fenofibrate, gallbladder investigations are indicated and this therapy should be discontinued (see sections 4.5 and 4.8).

### **Anticoagulants**

If Lipocomb is added to warfarin, another coumarin anticoagulant, or fluindione, the International Normalised Ratio (INR) should be appropriately monitored (see section 4.5).

### **Ciclosporin**

See sections 4.3 and 4.5.

### **Liver disease and alcohol**

Lipocomb should be used with caution in patients who consume excessive quantities of alcohol and/or have a history of liver disease.

### **Paediatric population**

The safety and efficacy of Lipocomb in children below the age of 18 years have not yet been established, therefore its use is not recommended in this age group.

## **4.5 Interaction with other medicines and other forms of interaction**

### **Contraindications**

#### **Ciclosporin**

During concomitant treatment with rosuvastatin and ciclosporin, rosuvastatin AUC values were on average 7 times higher than those observed in healthy volunteers (see section 4.3). Concomitant administration did not affect plasma concentrations of ciclosporin.

Co-administration of Lipocomb with ciclosporin is contraindicated (see section 4.3).

In a study of post-renal transplant patients with creatinine clearance of > 50 ml/min on a stable dose of ciclosporin, a single 10-mg dose of ezetimibe resulted in a 3,4-fold (range 2,3

to 7,9-fold) increase in the mean AUC for total ezetimibe compared to a healthy control population, receiving ezetimibe alone, from another study. In a different study, a renal transplant patient with severe renal impairment who was receiving ciclosporin and multiple other medications, demonstrated a 12 - fold greater exposure to total ezetimibe compared to concurrent controls receiving ezetimibe alone. In a study in healthy subjects, daily administration of 20 mg ezetimibe for 8 days with a single 100-mg dose of ciclosporin on Day 7 resulted in a mean 15 % increase in ciclosporin AUC (range 10 % decrease to 51 % increase) compared to a single 100 mg dose of ciclosporin alone. A controlled study on the effect of co-administered ezetimibe on ciclosporin exposure in renal transplant patients has not been conducted.

### **Combinations not recommended**

#### **Protease inhibitors**

Although the exact mechanism of interaction is unknown, concomitant protease inhibitor use may strongly increase rosuvastatin exposure (see section 4.5 Table). For instance, in a pharmacokinetic study, co-administration of 10 mg rosuvastatin and a combination product of two protease inhibitors (300 mg atazanavir / 100 mg ritonavir) in healthy volunteers was associated with an approximately three-fold and seven-fold increase in rosuvastatin AUC and C<sub>max</sub> respectively. The concomitant use of rosuvastatin and some protease inhibitor combinations may be considered after careful consideration of rosuvastatin dose adjustments based on the expected increase in rosuvastatin exposure (see sections 4.2, 4.4, and 4.5 Table). The combination is not suitable for initial therapy. Treatment initiation or dose adjustment if necessary should only be done with the monocomponents and after setting the appropriate doses the switch to the fixed dose combination of the appropriate strength is possible.

#### **Transporter protein inhibitors**

Rosuvastatin is a substrate for certain transporter proteins including the hepatic uptake transporter OATP1B1 and efflux transporter BCRP. Concomitant administration of

Lipocomb with medicines that are inhibitors of these transporter proteins may result in increased rosuvastatin plasma concentrations and an increased risk of myopathy (see sections 4.2, 4.4, and 4.5 Table).

### **Gemfibrozil and other lipid-lowering products**

Concomitant use of rosuvastatin and gemfibrozil resulted in a 2-fold increase in rosuvastatin C<sub>max</sub> and AUC (see section 4.4). Based on data from specific interaction studies no pharmacokinetic relevant interaction with fenofibrate is expected, however a pharmacodynamic interaction may occur.

Gemfibrozil, fenofibrate, other fibrates and lipid lowering doses (> or equal to 1 g/day) of niacin (nicotinic acid) increase the risk of myopathy when given concomitantly with HMG-CoA reductase inhibitors, probably because they can produce myopathy when given alone.

In patients receiving fenofibrate and ezetimibe, medical practitioners should be aware of the possible risk of cholelithiasis and gallbladder disease (see section 4.4 and 4.8). If cholelithiasis is suspected in a patient receiving ezetimibe and fenofibrate, gallbladder investigations are indicated

and this therapy should be discontinued (see section 4.8). Concomitant fenofibrate or gemfibrozil

administration modestly increased total ezetimibe concentrations (approximately 1,5- and 1,7-fold respectively). Co-administration of ezetimibe with other fibrates has not been studied. Fibrates may increase cholesterol excretion into the bile, leading to cholelithiasis.

In animal studies, ezetimibe sometimes increased cholesterol in the gallbladder bile, but not all species (see section 5.3). A lithogenic risk associated with the therapeutic use of ezetimibe cannot be ruled out.

### **Fusidic Acid**

The risk of myopathy including rhabdomyolysis may be increased by the concomitant administration of systemic fusidic acid with statins. The mechanism of this interaction (whether it is pharmacodynamic or pharmacokinetic, or both) is yet unknown. There have been reports of rhabdomyolysis (including some fatalities) in patients receiving this

combination.

If treatment with systemic fusidic acid is necessary, rosuvastatin treatment should be discontinued throughout the duration of the fusidic acid treatment. Also see section 4.4.

### **Other interactions**

#### **Antacid**

The simultaneous dosing of rosuvastatin with an antacid suspension containing aluminium and magnesium hydroxide resulted in a decrease in rosuvastatin plasma concentration of approximately 50 %. This effect was mitigated when the antacid was dosed 2 hours after rosuvastatin. The clinical relevance of this interaction has not been studied.

Concomitant antacid administration decreased the rate of absorption of ezetimibe but had no effect on the bioavailability of ezetimibe. This decreased rate of absorption is not considered clinically significant.

#### **Erythromycin**

Concomitant use of rosuvastatin and erythromycin resulted in a 20 % decrease in AUC<sub>0-t</sub> and a 30 % decrease in C<sub>max</sub> of rosuvastatin. This interaction may be caused by the increase in gut motility caused by erythromycin.

#### **Cytochrome P450 enzymes**

Results from in vitro and in vivo studies show that rosuvastatin is neither an inhibitor nor an inducer of cytochrome P450 isoenzymes. In addition, rosuvastatin is a poor substrate for these isoenzymes. Therefore, drug interactions resulting from cytochrome P450-mediated metabolism are not expected. No clinically relevant interactions have been observed between rosuvastatin and either fluconazole (an inhibitor of CYP2C9 and CYP3A4) or ketoconazole (an inhibitor of CYP2A6 and CYP3A4).

In preclinical studies, it has been shown that ezetimibe does not induce cytochrome P450 drug metabolising enzymes. No clinically significant pharmacokinetic interactions have been observed between ezetimibe and drugs known to be metabolised by cytochromes P450 1A2, 2D6, 2C8, 2C9, and 3A4, or N- acetyltransferase.

### **Vitamin K antagonists**

As with other HMG-CoA reductase inhibitors, the initiation of treatment or dosage up-titration of rosuvastatin in patients treated concomitantly with vitamin K antagonists (e.g. warfarin or another coumarin anticoagulant) may result in an increase in International Normalised Ratio (INR).

Discontinuation or down-titration of rosuvastatin may result in a decrease in INR. In such situations, appropriate monitoring of INR is desirable.

Concomitant administration of ezetimibe (10 mg once daily) had no effect on bioavailability of warfarin and prothrombin time in a study of twelve healthy adult males. However, there have been post-marketing reports of increased International Normalised Ratio (INR) in patients who had

ezetimibe added to warfarin or fluindione. If Lipocomb are added to warfarin, another coumarin

anticoagulant, or fluindione, INR should be appropriately monitored (see section 4.4).

### **Oral contraceptive/hormone replacement therapy (HRT)**

Concomitant use of rosuvastatin and oral contraceptive resulted in an increase in ethinyl oestradiol and norgestrel AUC of 26 % and 34 %, respectively. These increased plasma levels should be considered when selecting oral contraceptive doses. There are no pharmacokinetic data available in subjects taking concomitant rosuvastatin and HRT and therefore a similar effect cannot be excluded. However, the combination has been extensively used in women in clinical trials and was well tolerated.

In clinical interaction studies, ezetimibe had no effect on the pharmacokinetics of oral contraceptives (ethinyl oestradiol and levonorgestrel).

### **Colestyramine**

Concomitant colestyramine administration decreased the mean area under the curve (AUC) of total ezetimibe (ezetimibe + ezetimibe glucuronide) approximately 55 %. The incremental low-density lipoprotein cholesterol (LDL-C) reduction due to adding ezetimibe to colestyramine may be lessened by this interaction (see section 4.2).

## **Statins**

No clinically significant pharmacokinetic interactions were seen when ezetimibe was co-administered with atorvastatin, simvastatin, pravastatin, lovastatin, fluvastatin or rosuvastatin.

## **Other medicines**

Based on data from specific interaction studies no clinically relevant interaction between rosuvastatin and digoxin is expected.

In clinical interaction studies, ezetimibe had no effect on the pharmacokinetics of dapsone, dextromethorphan, digoxin, glipizide, tolbutamide, or midazolam, during co-administration. Cimetidine, co-administered with ezetimibe, had no effect on the bioavailability of ezetimibe.

## **Interactions requiring rosuvastatin dose adjustments (see also Table below)**

When it is necessary to co-administer rosuvastatin with other medicines known to increase exposure to rosuvastatin, doses should be adjusted. Start with a 5 mg once daily dose of rosuvastatin if the expected increase in exposure (AUC) is approximately 2-fold or higher. The maximum daily dose should be adjusted so that the expected rosuvastatin exposure would not likely exceed that of a 40 mg daily dose of rosuvastatin taken without interacting medicines, for example a 20 mg dose of rosuvastatin with gemfibrozil (1,9-fold increase), and a 10 mg dose of rosuvastatin with combination atazanavir/ritonavir (3,1-fold increase).

**Table 1 Effect of co-administered medicines on rosuvastatin exposure (AUC; in order of decreasing magnitude) from published clinical trials**

<b>Interacting drug dose regimen</b>	<b>Rosuvastatin dose regimen</b>	<b>Change in rosuvastatin AUC*</b>
Ciclosporin 75 mg BID to 200 mg BID, 6 months	10 mg OD, 10 days	7,1-fold ↑
Regorafenib 160 mg, OD, 14 days	5 mg single dose	3,8-fold ↑
Atazanavir 300 mg/ritonavir 100 mg OD, 8 days	10 mg, single dose	3,1-fold ↑
Simeprevir 150 mg OD, 7 days	10 mg single dose	2,8-fold ↑
Velpatasvir 100 mg OD	10 mg, single dose	2,7-fold ↑
Ombitasvir 25 mg/ paritaprevir 150 mg/ Ritonavir 100 mg OD/ dasabuvir 400 mg BID, 14 days	5 mg, single dose	2,6-fold ↑
Grazoprevir 200 mg/ elbasvir 50 mg OD, 11 days	10 mg, single dose	2,3-fold ↑
Glecaprevir 400 mg/ pibrentasvir 120 mg OD, 7 days	5 mg OD, 7 days	2,2-fold ↑
Lopinavir 400 mg/ritonavir 100 mg BID, 17 days	20 mg OD, 7 days	2,1-fold ↑
Clopidogrel 300 mg loading, followed by 75 mg at 24 hours	20 mg, single dose	2-fold ↑
Gemfibrozil 600 mg BID, 7 days	80 mg, single dose	1,9-fold ↑
Eltrombopag 75 mg OD, 5 days	10 mg, single dose	1,6-fold ↑
Darunavir 600 mg/ritonavir 100 mg BID, 7 days	10 mg OD, 7 days	1,5-fold ↑
Tipranavir 500 mg/ritonavir 200 mg BID, 11 days	10 mg, single dose	1,4-fold ↑
Dronedarone 400 mg BID	Not available	1,4-fold ↑
Itraconazole 200 mg OD, 5 days	10 mg, single dose	1,4-fold ↑**
Fosamprenavir 700 mg/ritonavir 100 mg BID, 8 days	10 mg, single dose	↔
Aleglitazar 0.3 mg, 7 days	40 mg, 7 days	↔
Silymarin 140 mg TID, 5 days	10 mg, single dose	↔

Fenofibrate 67 mg TID, 7 days	10 mg, 7 days	↔
Rifampin 450 mg OD, 7 days	20 mg, single dose	↔
Ketoconazole 200 mg BID, 7 days	80 mg, single dose	↔
Fluconazole 200 mg OD, 11 days	80 mg, single dose	↔
Erythromycin 500 mg QID, 7 days	80 mg, single dose	20 % ↓
Baicalin 50 mg TID, 14 days	20 mg, single dose	47 % ↓

\*Data given as x-fold change represent a simple ratio between co-administration and rosuvastatin alone. Data given as % change represent % difference relative to rosuvastatin alone.

Increase is indicated as “↑” no change as “↔” decrease as “↓”

\*\*Several interaction studies have been performed at different rosuvastatin dosages, the table shows the most significant ratio

OD = once daily; BID = twice daily; TID = three times daily; QID = four times daily.

The combination is not suitable for initial therapy. Treatment initiation or dose adjustment if necessary should only be done with the monocomponents and after setting the appropriate doses the switch to the fixed dose combination of the appropriate strength is possible.

#### **4.6 Fertility, pregnancy and lactation**

Lipocomb is contraindicated in pregnancy and breastfeeding.

Women of childbearing potential should use appropriate contraceptive measures.

##### **Pregnancy**

##### **Rosuvastatin**

Since cholesterol and other products of cholesterol biosynthesis are essential for the development of the foetus, the discontinuation of lipid-lowering medicines during pregnancy should have little impact on the outcome of long-term therapy of primary hypercholesterolaemia. Animal studies provide limited evidence of reproductive toxicity (see section 5.3). If a patient becomes pregnant during use of Lipocomb, treatment should be

discontinued immediately.

### **Ezetimibe**

No clinical data are available on the use of ezetimibe during pregnancy.

Animal studies on the use of ezetimibe in monotherapy have shown no evidence of direct or indirect harmful effects on pregnancy, embryofetal development, birth or postnatal development (see section 5.3). The use of ezetimibe is not recommended in pregnancy.

### **Breastfeeding**

### **Rosuvastatin**

Rosuvastatin is excreted in the milk of rats. There are no data with respect to excretion of rosuvastatin in milk in humans (see section 4.3).

### **Ezetimibe**

Studies on rats have shown that ezetimibe is secreted into milk. It is not known if ezetimibe is secreted into human breast milk.

### **Fertility**

No clinical trial data are available on the effects of ezetimibe on human fertility. Ezetimibe had no effect on the fertility of male or female rats (see section 5.3).

No studies on the effect on fertility have been conducted with rosuvastatin and ezetimibe in combination.

## **4.7 Effects on ability to drive and use machines**

Lipocomb has no or negligible influence on the ability to drive and use machines. Studies to determine the effect of rosuvastatin and/or ezetimibe on the ability to drive and use machines have not been conducted. However, when driving vehicles or operating machines, it should be taken into account that dizziness may occur during treatment.

## **4.8 Undesirable effects**

Summary of the safety profile.

According to available data 1 200 patients took the rosuvastatin and ezetimibe combination in clinical studies. As reported in the published literature, the most frequent common adverse events related to rosuvastatin- ezetimibe combination treatment in

hypercholesterolemic patients are increased hepatic transaminases, gastrointestinal problems and muscle pain. However, a pharmacodynamic interaction, in terms of adverse effects, between rosuvastatin and ezetimibe cannot be ruled out (see section 5.2).

Tabulated list of adverse reactions

<b>Organ class</b>	<b>Frequent</b>	<b>Less frequent</b>
<b>Blood and lymphatic system disorders</b>		Thrombocytopenia <sup>2, 5</sup>
<b>Immune system disorders</b>		Hypersensitivity reactions including angioedema <sup>2</sup> Hypersensitivity (including rash, urticaria, anaphylaxis and angioedema) <sup>5</sup>
<b>Endocrine disorders</b>	Diabetes mellitus <sup>1, 2</sup>	
<b>Metabolism and nutrition disorders</b>		Decreased appetite
<b>Psychiatric disorders</b>		Depression <sup>2, 5</sup>
<b>Nervous system disorders</b>	Headache <sup>2, 4</sup> Dizziness <sup>2</sup>	Paraesthesia <sup>4</sup> Polyneuropathy <sup>2</sup> Memory loss <sup>2</sup> Peripheral neuropathy <sup>2</sup> Sleep disturbances (including insomnia and nightmares) <sup>2</sup> Dizziness <sup>5</sup> Paraesthesia <sup>5</sup>
<b>Vascular disorders</b>		Hot flush <sup>3</sup> , Hypertension <sup>3</sup>
<b>Respiratory, thoracic and mediastinal disorders</b>		Cough <sup>2, 3</sup> Dyspnoea <sup>2, 5</sup>
<b>Gastrointestinal disorders</b>	Constipation <sup>2</sup> Nausea <sup>2</sup> Abdominal pain <sup>2, 3</sup> Diarrhoea <sup>3</sup> Flatulence <sup>3</sup>	Dyspepsia <sup>3</sup> Gastro-oesophageal reflux disease <sup>3</sup> Nausea <sup>3</sup> Dry mouth <sup>4</sup> Gastritis <sup>4</sup> Pancreatitis <sup>2, 5</sup>

		Diarrhoea <sup>2</sup> Constipation <sup>5</sup>
<b>Hepatobiliary disorders</b>		Increased hepatic transaminases <sup>2</sup> Jaundice <sup>2</sup> Hepatitis <sup>2, 5</sup> Cholelithiasis <sup>5</sup> Cholecystitis <sup>5</sup>
<b>Skin and subcutaneous tissue disorders</b>		Pruritus <sup>2, 4</sup> Rash <sup>2, 4</sup> Urticaria <sup>2, 4</sup> Stevens-Johnson syndrome <sup>2</sup> Erythema multiforme <sup>5</sup>
<b>Organ class</b>	<b>Frequent</b>	<b>Less frequent</b>

<b>Musculoskeletal and connective tissue disorders</b>	Myalgia <sup>2, 4</sup>	Arthralgia <sup>2, 3, 5</sup> Muscle spasms <sup>3</sup> Neck pain <sup>3</sup> Back pain <sup>4</sup> Muscular weakness <sup>4</sup> Pain in extremity <sup>4</sup> Myopathy (including myositis) <sup>2</sup> Rhabdomyolysis <sup>2</sup> Lupus like syndrome <sup>2</sup> Muscle rupture <sup>2</sup> Immune-mediated necrotising myopathy <sup>5</sup> Tendon disorders sometimes complicated by rupture <sup>2</sup> Myalgia <sup>5</sup> Myopathy/rhabdomyolysis <sup>5</sup> (see section 4.4)
<b>Renal and urinary disorders</b>		Haematuria <sup>2</sup>
<b>Reproductive system and breast disorders</b>		Gynecomastia <sup>2</sup>
<b>General disorders and administration site conditions</b>	Asthenia <sup>2</sup> , Fatigue <sup>3</sup>	Chest pain <sup>3</sup> Pain <sup>3</sup> Asthenia <sup>4, 5</sup> Peripheral oedema <sup>4</sup> Oedema <sup>2</sup>
<b>Investigations</b>	ALT and/or AST Increased <sup>4</sup>	ALT and/or AST increased <sup>3</sup> Blood CPK increased <sup>3</sup> Gamma- glutamyltransferase increased <sup>3</sup> Liver function test abnormal <sup>3</sup>

<sup>1</sup> Frequency will depend on the presence or absence of risk factors (fasting blood glucose  $\geq 5,6$  mmol/L, BMI  $> 30$  kg/m<sup>2</sup>, raised triglycerides, history of hypertension) – for rosuvastatin.

<sup>2</sup> Adverse reaction profile for rosuvastatin based on data from clinical studies and extensive post- marketing experience.

<sup>3</sup> Ezetimibe in monotherapy. Adverse reactions were observed in patients treated with ezetimibe (N=2 396) and at a greater incidence than placebo (N = 1 159).

<sup>4</sup> Ezetimibe co-administered with a statin. Adverse reactions were observed in patients with ezetimibe co- administered with a statin (N = 11 308) and at a greater incidence than statin administered alone (N = 9 361).

<sup>5</sup> Additional adverse reactions of ezetimibe, reported in post-marketing experience.

Because these adverse experiences have been identified from spontaneous reports, their true frequencies are not known and cannot be estimated.

As with other HMG-CoA reductase inhibitors, the incidence of adverse drug reactions tends to be dose dependent.

### **Renal effects**

Proteinuria, detected by dipstick testing and mostly tubular in origin, has been observed in patients treated with rosuvastatin. Shifts in urine protein from none or trace to ++ or more were seen in < 1 % of patients at some time during treatment with 10 and 20 mg, and in approximately 3 % of patients treated with 40 mg. A minor increase in shift from none or trace to + was observed with the 20 mg dose. In most cases, proteinuria decreases or disappears spontaneously on continued therapy. Review of data from clinical trials and post-marketing experience to date has not identified a causal association between proteinuria and acute or progressive renal disease.

Haematuria has been observed in patients treated with rosuvastatin and clinical trial data show that the occurrence is low.

### **Skeletal Muscle Effects**

Effects on skeletal muscle e.g. myalgia, myopathy (including myositis), and rhabdomyolysis with and without acute renal failure have been reported in rosuvastatin-treated patients with all doses and in particular with doses > 20 mg.

A dose-related increase in CK levels has been observed in patients taking rosuvastatin; the majority of cases were mild, asymptomatic and transient. If CK-levels are elevated ( $> 5 \times$  ULN), the treatment should be discontinued (see section 4.4).

### **Liver Effects**

As with other HMG-CoA reductase inhibitors, a dose-related increase in transaminases has been observed in a small number of patients taking rosuvastatin; the majority of cases were mild, asymptomatic and transient.

The following adverse events have been reported with some statins:

- Sexual dysfunction
- Exceptional cases of interstitial lung disease, especially with long term therapy (see section 4.4).

The reporting rates for rhabdomyolysis, serious renal events and serious hepatic events (consisting mainly of increased hepatic transaminases) are higher at the 40 mg rosuvastatin dose.

### **Laboratory values**

In controlled clinical monotherapy trials, the incidence of clinically important elevations in serum transaminases (ALT and/or AST  $\geq 3X$  ULN, consecutive) was similar between ezetimibe (0,5 %) and placebo (0,3 %). In co-administration trials, the incidence was 1,3 % for patients treated with ezetimibe co-administered with a statin and 0.4% for patients treated with a statin alone. These elevations were generally asymptomatic, not associated with cholestasis, and returned to baseline after discontinuation of therapy or with continued treatment (see section 4.4).

In clinical trials, CPK  $>10X$  ULN was reported for 4 of 1,674 (0,2 %) patients administered ezetimibe alone vs 1 of 786 (0,1 %) patients administered placebo, and for 1 of 917 (0,1 %) patients co-administered ezetimibe and a statin vs 4 of 929 (0,4 %) patients administered a statin alone. There was no excess of myopathy or rhabdomyolysis associated with ezetimibe compared with the relevant control arm (placebo or statin alone) (see section 4.4).

## **Paediatric population**

The safety and efficacy of Lipocomb in children below the age of 18 years have not yet been established (see section 5.1).

## **Rosuvastatin**

Creatine kinase elevations >10xULN and muscle symptoms following exercise or increased physical activity were observed more frequently in a 52-week clinical trial of children and adolescents compared to adults. In other respects, the safety profile of rosuvastatin was similar in children and adolescents compared to adults.

## **Ezetimibe**

### **Paediatric (6 to 17 years of age) patients**

In a study involving paediatric (6 to 10 years of age) patients with heterozygous familial or non-familial hypercholesterolaemia (n = 138), elevations of ALT and/or AST ( $\geq 3X$  ULN, consecutive) were observed in 1,1 % (1 patient) of the ezetimibe patients compared to 0 % in the placebo group. There were no elevations of CPK ( $\geq 10 X$  ULN). No cases of myopathy were reported.

In a separate study involving adolescent (10 to 17 years of age) patients with heterozygous familial hypercholesterolaemia (n = 248), elevations of ALT and/or AST ( $\geq 3 X$  ULN, consecutive) were observed in 3 % (4 patients) of the ezetimibe/simvastatin patients compared to 2 % (2 patients) in the simvastatin monotherapy group; these figures were respectively 2 % (2 patients) and 0 % for elevation of CPK ( $\geq 10 X$  ULN). No cases of myopathy were reported. These trials were not suited for comparison of rare adverse drug reactions.

## **Reporting of suspected adverse reactions**

Reporting suspected adverse reactions after authorisation of the medicine is important. It allows continued monitoring of the benefit/risk balance of the medicine. Healthcare professionals are asked to report any suspected adverse reactions to SAHPRA via the “**6.04**

**Adverse Drug Reaction Reporting Form**”, found online under SAHPRA’s publications:

<https://www.sahpra.org.za/Publications/Index/8>

## **4.9 Overdose**

There is no published literature data on rosuvastatin overdose.

There is no specific treatment in the event of overdose with rosuvastatin.

In clinical studies, administration of ezetimibe, 50 mg/day, to 15 healthy subjects for up to 14 days, or 40 mg/day to 18 patients with primary hypercholesterolaemia for up to 56 days, was generally well tolerated.

In the event of an overdose, symptomatic and supportive measures should be employed.

Liver

function and CK levels should be monitored. Haemodialysis is unlikely to be of benefit.

## **5. PHARMACOLOGICAL PROPERTIES**

### **5.1 Pharmacodynamic properties**

Pharmacotherapeutic group: lipid modifying agents; HMG CoA reductase inhibitors in combination with other lipid modifying agents, ATC code: C10BA06

#### **Rosuvastatin**

##### **Mechanism of action**

Rosuvastatin is a selective and competitive inhibitor of HMG-CoA reductase, the rate-limiting enzyme that converts 3-hydroxy-3-methylglutaryl coenzyme A to mevalonate, a precursor for cholesterol. The primary site of action of rosuvastatin is the liver, the target organ for cholesterol lowering.

Rosuvastatin increases the number of hepatic LDL receptors on the cell-surface, enhancing uptake and catabolism of LDL and it inhibits the hepatic synthesis of VLDL, thereby reducing the total number of VLDL and LDL particles.

##### **Pharmacodynamic effects**

Rosuvastatin reduces elevated LDL-cholesterol, total cholesterol and triglycerides and increases HDL-cholesterol. It also lowers ApoB, nonHDL-C, VLDL-C, VLDL-TG and increases ApoA-I (see Table below). Rosuvastatin also lowers the LDL-C/HDL-C, total C/HDL-C and nonHDL-C/HDL-C and the ApoB/ApoA-I ratios.

Dose response in patients with primary hypercholesterolaemia (type IIa and IIb)

(adjusted mean percent change from baseline)

Dose	N	LDL-C	Total-C	HDL-C	TG	nonHDL-C	ApoB	ApoA-I
Placebo	13	- 7	- 5	3	- 3	- 7	- 3	0
5 mg	17	- 45	- 33	13	- 35	- 44	- 38	4
10 mg	17	- 52	- 36	14	- 10	- 48	- 42	4
20 mg	17	- 55	- 40	8	- 23	- 51	- 46	5
40 mg	18	- 63	- 46	10	- 28	- 60	- 54	0

A therapeutic effect is obtained within 1 week following treatment initiation and 90 % of maximum response is achieved in 2 weeks. The maximum response is usually achieved by 4 weeks and is maintained after that.

### **Ezetimibe**

#### **Mechanism of action**

Ezetimibe is in a new class of lipid-lowering compounds that selectively inhibit the intestinal absorption of cholesterol and related plant sterols. Ezetimibe is orally active, and has a mechanism of action that differs from other classes of cholesterol-reducing compounds (e.g. statins, bile acid sequestrants [resins], fibric acid derivatives, and plant stanols). The molecular target of ezetimibe is the sterol transporter, Niemann-Pick C1-Like 1 (NPC1L1), which is responsible for the intestinal uptake of cholesterol and phytosterols. Ezetimibe localises at the brush border of the small intestine and inhibits the absorption of cholesterol, leading to a decrease in the delivery of intestinal cholesterol to the liver; statins reduce cholesterol synthesis in the liver and together these distinct mechanisms provide complementary cholesterol reduction. In a 2-week clinical study in 18 hypercholesterolaemic patients, ezetimibe inhibited intestinal cholesterol absorption by 54 %, compared with placebo.

#### **Pharmacodynamic effects**

A series of preclinical studies was performed to determine the selectivity of ezetimibe for inhibiting cholesterol absorption. Ezetimibe inhibited the absorption of [<sup>14</sup>C]-cholesterol with no effect on the absorption of triglycerides, fatty acids, bile acids, progesterone, ethinyl estradiol, or fat soluble vitamins A and D.

Epidemiologic studies have established that cardiovascular morbidity and mortality vary directly with the level of total-C and LDL-C and inversely with the level of HDL-C. A beneficial effect of ezetimibe on cardiovascular morbidity and mortality has not yet been demonstrated.

### **Rosuvastatin-ezetimibe co-administration**

#### **Clinical efficacy and safety**

A 6-week, randomised, double-blind, parallel-group, clinical trial evaluated the safety and efficacy of ezetimibe (10 mg) added to stable rosuvastatin therapy versus up-titration of rosuvastatin from 5 to 10 mg or from 10 to 20 mg (n = 440). Pooled data demonstrated that ezetimibe added to stable rosuvastatin 5 mg or 10 mg reduced LDL cholesterol by 21 %. In contrast, doubling rosuvastatin to

10 mg or 20 mg reduced LDL cholesterol by 5,7 % (between-group difference of 15,2 %, p < 0,001). Individually, ezetimibe plus rosuvastatin 5 mg reduced LDL cholesterol more than did rosuvastatin 10 mg (12,3 % difference, p < 0,001), and ezetimibe plus rosuvastatin 10 mg reduced LDL cholesterol more than did rosuvastatin 20 mg (17,5 % difference, p < 0,001).

A 6-week, randomised study was designed to investigate the efficacy and safety of rosuvastatin 40 mg alone or in combination with ezetimibe 10 mg in patients at high risk of coronary heart disease (n = 469). Significantly more patients receiving rosuvastatin/ezetimibe than rosuvastatin alone achieved their ATP III LDL cholesterol goal (< 100 mg/dl, 94,0 % vs 79,1 %, p < 0,001). Rosuvastatin 40 mg was effective at improving the atherogenic lipid profile in this high-risk population.

A randomised, open-label, 12-week study investigated the level of LDL reduction in each treatment arm (rosuvastatin 10 mg plus ezetimibe 10 mg, rosuvastatin 20 mg/ezetimibe 10 mg, simvastatin 40/ezetimibe 10 mg, simvastatin 80/ezetimibe 10 mg). The reduction from baseline with the low dose rosuvastatin combinations was 59,7 %, significantly superior to the low dose simvastatin combinations, 55,2 % (p < 0,05). Treatment with the high-dose rosuvastatin combo reduced LDL cholesterol 63,5 % compared with a reduction of 57,4 % with the high-dose simvastatin combination (p < 0,001).

## **Paediatric population**

See section 4.2 for information on paediatric use.

## **5.2 Pharmacokinetic properties**

### **Rosuvastatin and ezetimibe combination therapy**

Concomitant use of 10 mg rosuvastatin and 10 mg ezetimibe resulted in a 1,2 fold increase in AUC of rosuvastatin in hypercholesterolaemic subjects. A pharmacodynamic interaction, in terms of adverse effects, between rosuvastatin and ezetimibe cannot be ruled out.

### **Rosuvastatin**

#### **Absorption**

Maximum rosuvastatin plasma concentrations are achieved approximately 5 hours after oral administration. The absolute bioavailability is approximately 20 %.

#### **Distribution**

Rosuvastatin is taken up extensively by the liver, which is the primary site of cholesterol synthesis and LDL-C clearance. The volume of distribution of rosuvastatin is approximately 134 L. Approximately 90 % of rosuvastatin is bound to plasma proteins, mainly to albumin.

#### **Biotransformation**

Rosuvastatin undergoes limited metabolism (approximately 10 %). In vitro metabolism studies using human hepatocytes indicate that rosuvastatin is a poor substrate for cytochrome P450-based metabolism. CYP2C9 was the principal isoenzyme involved, with 2C19, 3A4 and 2D6 involved to a lesser extent. The main metabolites identified are the N-desmethyl- and lactone metabolites. The N-desmethyl metabolite is approximately 50 % less active than rosuvastatin whereas the lactone form is considered clinically inactive. Rosuvastatin accounts for greater than 90 % of the circulating HMG-CoA reductase inhibitor activity.

#### **Elimination**

Approximately 90 % of the rosuvastatin dose is excreted unchanged in the faeces (consisting of absorbed and non-absorbed active substance) and the remaining part is

excreted in urine.

Approximately 5 % is excreted unchanged in urine. The plasma elimination half-life is approximately 19 hours. The elimination half-life does not increase at higher doses. The geometric mean plasma clearance is approximately 50 litres/hour (coefficient of variation 21,7 %).

As with other HMG-CoA reductase inhibitors, the hepatic uptake of rosuvastatin involves the membrane transporter OATP-C. This transporter is important in the hepatic elimination of rosuvastatin.

### **Linearity**

Systemic exposure of rosuvastatin increases in proportion to dose. There are no changes in pharmacokinetic parameters following multiple daily doses.

### **Special populations**

#### **Age and Gender**

There was no clinically relevant effect of age or gender on the pharmacokinetics of rosuvastatin in adults. The pharmacokinetics of rosuvastatin in children and adolescents with heterozygous familial hypercholesterolaemia was similar to that of adult volunteers (see "Paediatric population" below).

#### **Race**

Pharmacokinetic studies show an approximate 2-fold elevation in median AUC and C<sub>max</sub> in Asian subjects (Japanese, Chinese, Filipino, Vietnamese and Koreans) compared with Caucasians; Asian-Indians show an approximate 1,3-fold elevation in median AUC and C<sub>max</sub>.

A population pharmacokinetic analysis revealed no clinically relevant differences in pharmacokinetics between Caucasian and Black groups.

#### **Renal impairment**

In a study in subjects with varying degrees of renal impairment, mild to moderate renal disease had no influence on plasma concentration of rosuvastatin or the N-desmethyl

metabolite. Subjects with severe impairment (CrCl < 30 ml/min) had a 3-fold increase in plasma concentration and a 9-fold increase in the N-desmethyl metabolite concentration compared to healthy volunteers. Steady-state plasma concentrations of rosuvastatin in subjects undergoing haemodialysis were 50 % greater compared to healthy volunteers.

### **Hepatic impairment**

In a study with subjects with varying degrees of hepatic impairment there was no evidence of increased exposure to rosuvastatin in subjects with Child-Pugh scores of 7 or below.

However, two subjects with Child-Pugh scores of 8 and 9 showed an increase in systemic exposure

of at least 2-fold compared to subjects with lower Child-Pugh scores.

There is no experience in subjects with Child-Pugh scores above 9.

### **Genetic polymorphisms**

Disposition of HMG-CoA reductase inhibitors, including rosuvastatin, involves OATP1B1 and BCRP transporter proteins. In patients with SLCO1B1 (OATP1B1) and/or ABCG2 (BCRP) genetic polymorphisms there is a risk of increased rosuvastatin exposure.

Individual polymorphisms of SLCO1B1 c.521CC and ABCG2 c.421AA are associated with a higher rosuvastatin exposure (AUC) compared to the SLCO1B1 c.521TT or ABCG2 c.421CC genotypes. This specific genotyping is not established in clinical practice, but for patients who are known to have these types of polymorphisms, a lower daily dose of Lipocomb is recommended.

### **Paediatric population**

Two pharmacokinetic studies with rosuvastatin (given as tablets) in paediatric patients with heterozygous familial hypercholesterolaemia 10-17 or 6-17 years of age (total of 214 patients) demonstrated that exposure in paediatric patients appears comparable to or lower than that in adult patients. Rosuvastatin exposure was predictable with respect to dose and time over a 2-year period.

## **Ezetimibe**

### **Absorption**

After oral administration, ezetimibe is rapidly absorbed and extensively conjugated to a pharmacologically-active phenolic glucuronide (ezetimibe-glucuronide). Mean maximum plasma concentrations (C<sub>max</sub>) occur within 1 to 2 hours for ezetimibe-glucuronide and 4 to 12 hours for ezetimibe. The absolute bioavailability of ezetimibe cannot be determined as the compound is virtually insoluble in aqueous media suitable for injection.

Concomitant food administration (high fat or non-fat meals) had no effect on the oral bioavailability of ezetimibe. Ezetimibe can be administered with or without food.

### **Distribution**

Ezetimibe and ezetimibe-glucuronide are bound 99,7 % and 88 to 92 % to human plasma proteins, respectively.

### **Biotransformation**

Ezetimibe is metabolised primarily in the small intestine and liver via glucuronide conjugation (a phase II reaction) with subsequent biliary excretion. Minimal oxidative metabolism (a phase I reaction) has been observed in all species evaluated. Ezetimibe and ezetimibe-glucuronide are the major drug-derived compounds detected in plasma, constituting approximately 10 to 20 % and 80 to 90 % of the total drug in plasma, respectively. Both ezetimibe and ezetimibe-glucuronide are slowly eliminated from plasma with evidence of significant enterohepatic recycling. The half-life for ezetimibe and ezetimibe-glucuronide is approximately 22 hours.

### **Elimination**

Following oral administration of <sup>14</sup>C-ezetimibe (20 mg) to human subjects, total ezetimibe accounted for approximately 93 % of the total radioactivity in plasma. Approximately 78 % and 11 % of the administered radioactivity were recovered in the faeces and urine, respectively, over a 10-day collection period. After 48 hours, there were no detectable levels of radioactivity in the plasma.

## **Special populations**

### **Age and Gender**

Plasma concentrations for total ezetimibe are about 2-fold higher in the elderly ( $\geq 65$  years) than in the young (18 to 45 years). LDL-C reduction and safety profile are comparable between elderly and young subjects treated with ezetimibe. Therefore, no dosage adjustment is necessary in the elderly.

Plasma concentrations for total ezetimibe are slightly higher (approximately 20 %) in women than in men. LDL-C reduction and safety profile are comparable between men and women treated with ezetimibe.

Therefore, no dosage adjustment is necessary on the basis of gender.

### **Renal impairment**

After a single 10 mg dose of ezetimibe in patients with severe renal disease ( $n = 8$ ; mean  $\text{CrCl} \leq 30 \text{ ml/min/1.73 m}^2$ ), the mean AUC for total ezetimibe was increased approximately 1.5-fold, compared to healthy subjects ( $n = 9$ ). This result is not considered clinically significant. No dosage adjustment is necessary for renally impaired patients.

An additional patient in this study (post-renal transplant and receiving multiple medications, including ciclosporin) had a 12-fold greater exposure to total ezetimibe.

### **Hepatic impairment**

After a single 10 mg dose of ezetimibe, the mean AUC for total ezetimibe was increased approximately 1.7-fold in patients with mild hepatic impairment (Child Pugh score 5 or 6), compared to healthy subjects. In a 14-day, multiple-dose study (10 mg daily) in patients with moderate hepatic impairment (Child Pugh score 7 to 9), the mean AUC for total ezetimibe was increased approximately 4-fold on Day 1 and Day 14 compared to healthy subjects. No dosage adjustment is necessary for patients with mild hepatic impairment.

Due to the unknown effects of the increased exposure to ezetimibe in patients with moderate or severe (Child Pugh score  $> 9$ ) hepatic impairment, Lipocomb is not recommended in these patients (see section 4.4).

### **Paediatric population**

The pharmacokinetics of ezetimibe are similar between children  $\geq 6$  years and adults. Pharmacokinetic data in the paediatric population  $< 6$  years of age are not available. Clinical experience in paediatric and adolescent patients includes patients with HoFH, HeFH, or sitosterolaemia.

### **5.3 Preclinical safety data**

In co-administration studies with ezetimibe and statins the toxic effects observed were essentially those typically associated with statins. Some of the toxic effects were more pronounced than observed during treatment with statins alone. This is attributed to pharmacokinetic and pharmacodynamic interactions in co-administration therapy. No such interactions occurred in the clinical studies. Myopathies occurred in rats only after exposure to doses that were several times higher than the human therapeutic dose (approximately 20 times the AUC level for statins and 500 to 2 000 times the AUC level for the active metabolites).

In a series of in vivo and in vitro assays ezetimibe, given alone or co-administered with statins, exhibited no genotoxic potential. Long-term carcinogenicity tests on ezetimibe were negative.

The co-administration of ezetimibe and statins was not teratogenic in rats. In pregnant rabbits a small number of skeletal deformities (fused thoracic and caudal vertebrae, reduced number of caudal vertebrae) were observed.

### **Rosuvastatin**

Preclinical data reveal no special hazard for humans based on conventional studies of safety

pharmacology, genotoxicity and carcinogenicity potential. Specific tests for effects on hERG have not been evaluated. Adverse reactions not observed in clinical studies, but seen in animals at exposure levels similar to clinical exposure levels were as follows: in repeated-dose toxicity studies histopathologic liver changes likely due to the pharmacologic action of rosuvastatin were observed in mouse, rat, and to a lesser extent with effects in the gall bladder in dogs, but not in monkeys. In addition, testicular toxicity was observed in

monkeys and dogs at higher dosages. Reproductive toxicity was evident in rats, with reduced litter sizes, litter weight and pup survival observed at maternally toxic doses, where systemic exposures were several times above the therapeutic exposure level.

### **Ezetimibe**

Animal studies on the chronic toxicity of ezetimibe identified no target organs for toxic effects. In dogs treated for four weeks with ezetimibe ( $\geq 0,03$  mg/kg/day) the cholesterol concentration in the cystic bile was increased by a factor of 2,5 to 3,5. However, in a one-year study on dogs given doses of up to 300 mg/kg/day no increased incidence of cholelithiasis or other hepatobiliary effects were observed.

The significance of these data for humans is not known. A lithogenic risk associated with the therapeutic use of ezetimibe cannot be ruled out.

Ezetimibe had no effect on the fertility of male or female rats, nor was it found to be teratogenic in

rats or rabbits, nor did it affect prenatal or postnatal development. Ezetimibe crossed the placental barrier in pregnant rats and rabbits given multiple doses of 1 000 mg/kg/day. The co-administration of ezetimibe with lovastatin resulted in embryo lethal effects.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

#### **Core**

Silicified microcrystalline cellulose (Microcrystalline Cellulose (E460) and Colloidal Anhydrous Silica (E551) Colloidal Anhydrous Silica (E551), Magnesium stearate (E572), Povidone (E1201)

Croscarmellose Sodium (E468), Microcrystalline Cellulose (E460), Mannitol (E421), Sodium laurilsulfate (E514), Low-substituted hydroxypropyl cellulose (E463)

#### **Capsule shell**

#### **Lipocomb 10 mg/10 mg hard gelatin capsules**

Cap and body: Yellow iron oxide (E172), Titanium dioxide (E171), Gelatin

#### **Lipocomb 20 mg/10 mg hard gelatin capsules**

Cap: Red iron oxide (E172), Titanium dioxide (E171), Yellow iron oxide (E172),

Gelatin Body: Yellow iron oxide (E172), Titanium dioxide (E171), Gelatine

## **6.2 Incompatibilities**

Not applicable.

## **6.3 Shelf life**

3 years

## **6.4 Special precautions for storage**

Store at or below 25 °C.

Keep out of reach of children.

Store in the original package in order to protect from light and moisture.

## **6.5 Nature and contents of container**

Packs of 30 hard gelatin capsules in cold blister (OPA/AL/PVC//Al) in a carton.

## **6.6 Special precautions for disposal and other handling**

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

## **7. HOLDER OF CERTIFICATE OF REGISTRATION**

Servier Laboratories SA (Pty) Ltd

3rd Floor, Building J

Hertford Office Park

90 Bekker Road

Vorna Valley

Midrand

2196

## **8. REGISTRATION NUMBER(S)**

Lipocomb: 10 mg/ 10 mg: 55/7.5/0407

Lipocomb: 20 mg/ 10 mg: 55/7.5/0408

**9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

9 May 2023

**10. DATE OF REVISION OF THE TEXT**