

**Applicant/PHCR:** AUROGEN SA (PTY) LTD

**Product proprietary name:** MOLNARZ

**Dosage form and strength:** Hard capsules; 200 mg

**Submitted:** 17/05/2023

### 1.3.1.1 PROFESSIONAL INFORMATION FOR MEDICINES FOR HUMAN USE

(APPROVED COPY)

#### SCHEDULING STATUS

**S4**

#### 1. NAME OF THE MEDICINE

**MOLNARZ**, 200 mg (hard capsules)

#### 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each hard capsule contains 200 mg of molnupiravir. For the full list of excipients, see section

6.1.

Sugar free.

For a full list of excipients, see section 6.1.

#### 3. PHARMACEUTICAL FORM

**MOLNARZ** (hard capsules)

White to off white granular powder filled in size '0' hard capsule with green opaque coloured cap and body imprinted with "MOL" on cap and '200' on body with BLACK TEK ink.

#### 4. CLINICAL PARTICULARS

##### 4.1. Therapeutic indications

**MOLNARZ** is indicated for treatment of mild to moderate coronavirus disease 2019 (COVID-19) in adults with a positive SARS-COV-2 diagnostic test and who do not require

**Applicant/PHCR:** AUROGEN SA (PTY) LTD

**Product proprietary name:** MOLNARZ

**Dosage form and strength:** Hard capsules; 200 mg

**Submitted:** 17/05/2023

supplemental oxygen due to COVID-19 and who have at least one risk factor for developing severe illness (see 5.1 for clinical studies).

#### **4.2. Posology and method of administration**

##### **Posology**

###### *Adults*

The recommended dose of **MOLNARZ** in adult patients is 800 mg (four 200 mg capsules) taken orally every 12 hours for 5 days, with or without food. Should a patient require hospitalization after starting treatment with **MOLNARZ**, the patient may complete the full 5 day treatment course per the healthcare provider's discretion.

The safety and efficacy of **MOLNARZ** when administered for periods longer than 5 days have not been established.

**MOLNARZ** should be administered as soon as possible after a diagnosis of COVID-19 has been made and within 5 days of symptom onset in adults who are at risk for progression to severe COVID-19, including hospitalization or death. Certain medical conditions or other factors may place individual patients at increased risk for progression to severe COVID-19 [see section 5.1 Clinical Studies].

###### *Missed dose*

If the patient misses a dose of **MOLNARZ** within 10 hours of the time it is usually taken, the patient should take as soon as possible and resume the normal dosing schedule. If a patient misses a dose by more than 10 hours, the patient should not take the missed dose and instead take the next dose at the regularly scheduled time. The patient should not double the dose to make up for a missed dose.

##### **Special populations**

**Applicant/PHCR:** AUROGEN SA (PTY) LTD  
**Product proprietary name:** MOLNARZ  
**Dosage form and strength:** Hard capsules; 200 mg

**Submitted:** 17/05/2023

#### *Elderly*

No dose adjustment of **MOLNARZ** is recommended for elderly patients (see section 5.2 Gender, Race and Age).

#### *Renal impairment*

The pharmacokinetics of molnupiravir and NHC has not been evaluated in patients with eGFR less than 30 mL/min or on dialysis.

#### *Hepatic impairment*

No dose adjustment of **MOLNARZ** is recommended for patients with hepatic impairment (see section 5.2 Hepatic Impairment).

#### *Paediatric population*

The safety and efficacy of **MOLNARZ** in patients below 18 years of age have not been established. No data are available (see sections 5.2 Paediatric Population and 5.3 General Toxicity).

#### *Pregnancy*

Based on animal data, **MOLNARZ** may cause foetal harm. Human pregnancy data are not available. The use of **MOLNARZ** is not recommended during pregnancy (see section 5.3 Development).

#### **Method of administration**

For oral use.

**MOLNARZ** can be taken with or without food.

**Applicant/PHCR:** AUROGEN SA (PTY) LTD

**Product proprietary name:** MOLNARZ

**Dosage form and strength:** Hard capsules; 200 mg

**Submitted:** 17/05/2023

The capsules should be swallowed whole with a sufficient amount of fluid (e.g., a glass of water). The capsules should not be opened, crushed or chewed.

#### **4.3. Contraindications**

**MOLNARZ** is contraindicated in patients who have a known history of hypersensitivity to the active substance or any of the other ingredients (see Section 6.1).

#### **4.4. Special warnings and precautions for use**

##### **Sodium**

This medicinal product contains less than 1 mmol sodium (23 mg) per dose of 4 capsules, that is to say essentially 'sodium-free'.

#### **4.5 Interaction with other medicines and other forms of interaction**

No interactions have been identified based on the limited available data. No clinical interaction studies have been performed with molnupiravir. Molnupiravir is hydrolysed to n-hydroxycytidine (NHC) prior to reaching systemic circulation. Uptake of NHC and metabolism to NHC-TP are mediated by the same pathways involved in endogenous pyrimidine metabolism. NHC is not a substrate of major medicine metabolising enzymes or transporters. Based on *in vitro* studies, neither molnupiravir nor NHC are inhibitors or inducers of major medicine metabolising enzymes or inhibitors of major medicine transporters. Therefore, the potential for molnupiravir or NHC to interact with concomitant medications is considered unlikely.

#### **4.6 Fertility, pregnancy and lactation**

**Applicant/PHCR:** AUROGEN SA (PTY) LTD  
**Product proprietary name:** MOLNARZ  
**Dosage form and strength:** Hard capsules; 200 mg

**Submitted:** 17/05/2023

#### Women of childbearing potential

Advise women of childbearing potential to use effective contraception for the duration of treatment and for 4 days after the last dose of **MOLNARZ**.

#### Pregnancy

##### Risk Summary

Based on animal data, **MOLNARZ** may cause foetal harm when administered to pregnant women. There are no available data on the use of **MOLNARZ** in pregnant women to evaluate the risk of major birth defects, miscarriage or adverse maternal or foetal outcomes. The use of **MOLNARZ** is not recommended during pregnancy. In an animal reproduction study, oral administration of molnupiravir to pregnant rats during the period of organogenesis resulted in embryofoetal lethality and teratogenicity at 8 times the human NHC exposures at the recommended human dose (RHD) and reduced foetal growth at  $\geq 3$  times the human NHC exposure at the RHD. Oral administration of molnupiravir to rabbits during the period of organogenesis resulted in reduced foetal body weights at 18 times the human NHC exposure at the RHD [see section 5.3 Development]. In a pre- and post-natal developmental study, molnupiravir was administered orally to female rats at doses up to 500 mg/kg/day (1.6 times the human NHC exposure at the RHD) from GD6 through lactation day 20. No effects were observed in offspring.

#### **Breastfeeding**

It is unknown whether molnupiravir or any of the components of molnupiravir are present in human milk, affect human milk production, or have effect on the breastfed infant. Animal lactation studies with molnupiravir have not been conducted was detected in the plasma of nursing pups from lactating rats administered molnupiravir.

**Applicant/PHCR:** AUROGEN SA (PTY) LTD  
**Product proprietary name:** MOLNARZ  
**Dosage form and strength:** Hard capsules; 200 mg

**Submitted:** 17/05/2023

Based on the potential for adverse reactions on the infant from **MOLNARZ**, breastfeeding is not recommended during treatment and for 4 days after the last dose of **MOLNARZ**.

## Fertility

There were no effects on female or male fertility in rats at NHC exposures approximately 2 and 6 times respectively, the exposure in humans at the recommended human dose (RHD).

## 4.7 Effects on ability to drive and use machines

**MOLNARZ** has no or negligible influence on the ability to drive and use machines.

## 4.8 Undesirable effects

### a) Summary of safety profile

The most common adverse reactions in patients treated with 800 mg Molnupiravir every 12 hours for 5 days in the Phase 3 MOVE-OUT clinical trial were diarrhoea (2 %), nausea (1 %), and dizziness (1 %) all of which were Grade 1 (mild) or Grade 2 (moderate) in severity.

### Tabulated summary of adverse reactions

The adverse reactions are listed below by MedDRA system organ class and frequency. Frequencies are defined as follows: Frequent = very common and common; Less frequent = rare and very rare; not known (cannot be estimated from the available data).

**Table 1: Tabulated summary of adverse reactions**

Adverse Reaction	Frequency
<i>Immune systems disorders</i>	
Hypersensitivity	Less frequent

**Applicant/PHCR:** AUROGEN SA (PTY) LTD

**Product proprietary name:** MOLNARZ

**Dosage form and strength:** Hard capsules; 200 mg

**Submitted:** 17/05/2023

<i>Nervous system disorders</i>	
dizziness	Frequent
headache	Frequent
<i>Gastrointestinal disorders</i>	
diarrhoea, nausea	Frequent
vomiting	Less frequent
<i>Skin and subcutaneous tissue disorders</i>	
Angioedema, erythema, rash, urticaria	Less frequent

### **Reporting of suspected adverse reactions**

Reporting suspected adverse reactions after authorisation of the medicine is important. It allows continued monitoring of the benefit/risk balance of the medicine. Healthcare providers are asked to report any suspected adverse reactions to SAHPRA via the “**6.04 Adverse Medicine Reactions Reporting Form**”, found online under SAHPRA’s publications:

<https://www.sahpra.org.za/Publications/Index/8>.

### **4.9 Overdose**

There is no human experience of overdosage with **MOLNARZ**. Treatment of overdose with **MOLNARZ** should consist of general supportive measures including the monitoring of the clinical status of the patient. Haemodialysis is not expected to result in effective elimination of NHC.

## **5 PHARMACOLOGICAL PROPERTIES**

### **5.1 Pharmacodynamic properties**

A20.2.8 Antiviral agents

Pharmacotherapeutic group: Antiviral for systemic use, direct acting antivirals

**Applicant/PHCR:** AUROGEN SA (PTY) LTD

**Product proprietary name:** MOLNARZ

**Dosage form and strength:** Hard capsules; 200 mg

**Submitted:** 17/05/2023

The relationship between NHC and intracellular NHC-TP with antiviral efficacy has not been evaluated clinically.

#### Mechanism of action

Molnupiravir is a prodrug that is metabolised to the ribonucleoside analogue N-hydroxycytidine (NHC) which distributes into cells where it is phosphorylated to form the pharmacologically active ribonucleoside triphosphate (NHC-TP). NHC-TP acts by a mechanism known as viral error catastrophe. NHC-TP incorporation into viral RNA by the viral RNA polymerase, results in an accumulation of errors in the viral genome leading to inhibition of replication.

#### Microbiology

##### Antiviral Activity

NHC was active in cell culture assays against SARS-CoV-2 with 50% effective concentrations (EC<sub>50</sub>) ranging between 0.67 to 2.66 µM in A-549 cells and 0.32 to 2.03 µM in Vero E6 cells. NHC had similar activity against SARS-CoV-2 variants B.1.1.7 (Alpha), B.1351 (Beta), P.1 (Gamma), and B.1.617.2 (Delta) with EC<sub>50</sub> values of 1.59, 1.77 and 1.32 and 1.68 µM, respectively. No impact was observed on the *in vitro* antiviral activity of NHC against SARS-CoV-2 when NHC was tested in combination with abacavir, emtricitabine, hydroxychloroquine, lamivudine, nelfinavir, remdesivir, ribavirin, sofosbuvir, or tenofovir.

##### *Resistance*

No amino acid substitutions in SARS-CoV-2 associated with resistance to NHC have been identified in Phase 2 clinical trials evaluating molnupiravir for the treatment of COVID-19. Studies to evaluate selection of resistance to NHC with SARS-CoV-2 in cell culture have not been completed. Resistance selection studies have been conducted with other coronaviruses

**Applicant/PHCR: AUROGEN SA (PTY) LTD**

**Product proprietary name: MOLNARZ**

**Dosage form and strength: Hard capsules; 200 mg**

**Submitted: 17/05/2023**

(MHV and MERS-CoV) and showed a low likelihood of resistance development to NHC. Following 30 passages in cell culture, only a 2-fold decrease in susceptibility was observed and no NHC resistance associated amino acid substitutions were identified. NHC retained activity *in vitro* against virus with polymerase substitutions (e.g., F480L, V557L and E802D) associated with decreased remdesivir sensitivity, indicating a lack of cross-resistance.

#### *Activity against SARS-CoV-2 in animal models*

The antiviral activity of molnupiravir has been demonstrated in mouse, hamster and ferret models of SARS-CoV-2 infection. In mice, molnupiravir significantly reduced infectious SARS-CoV-2 levels in infected transplanted human lung tissue. In SARSCoV-2 infected ferrets, molnupiravir significantly reduced SARS-CoV-2 viral titers in the upper respiratory tract and completely inhibited viral spread to untreated contact animals. In SARS-CoV-2 infected Syrian hamsters, molnupiravir reduced viral RNA and infectious virus titers in the lungs of animals. Histopathological analysis of lung tissue harvested after infection, showed significantly reduced SARS-CoV-2 viral antigen levels and a lower abundance of pulmonary lesions in molnupiravir-treated animals compared with controls.

#### Clinical Studies

Clinical data are based on data from 1,433 randomized subjects in the Phase 3 MOVE- OUT trial. MOVE-OUT is a randomized, placebo-controlled, double-blind clinical trial studying molnupiravir for the treatment of non-hospitalized patients with mild to moderate COVID-19 who are at risk for progressing to severe COVID-19 and/or hospitalization. Eligible subjects were 18 years of age and older and had one or more pre-defined risk factors for disease progression: over 60 years of age, diabetes, obesity (BMI  $\geq 30$ ), chronic kidney disease, serious heart conditions, chronic obstructive pulmonary disease, or active cancer. The study included symptomatic subjects not vaccinated against SARS-CoV-2 and who had laboratory

**Applicant/PHCR:** AUROGEN SA (PTY) LTD

**Product proprietary name:** MOLNARZ

**Dosage form and strength:** Hard capsules; 200 mg

**Submitted:** 17/05/2023

confirmed SARS-CoV-2 infection and symptom onset within 5 days of randomization.

Subjects were randomized 1:1 to receive 800 mg of molnupiravir or placebo orally twice daily for 5 days.

At baseline, in all randomized subjects, the median age was 43 years (range: 18 to 90); 17 % of subjects were over 60 years of age and 3 % were 75 years of age or older; 49 % of subjects were male; 57 % were White, 5 % Black or African American, 3 % Asian, 50 % Hispanic or Latino. Forty-eight percent of subjects received Molnupiravir or placebo within 3 days of COVID-19 symptom onset. The most common risk factors were obesity (74 %), over 60 years of age (17 %), and diabetes (16 %). Overall, baseline demographic and disease characteristics were well balanced between the treatment arms.

Table 2 provides the results of the primary endpoint (the percentage of subjects who were hospitalized or died through Day 29 due to any cause). Please refer to Figure 1 for results by certain subgroups.

**Table 2. Interim Efficacy Results in Non-Hospitalized Adults with COVID-19**

<b>Molnupiravir (N=709)</b> <b>n (%)</b>	<b>Placebo (N=699)</b> <b>n (%)</b>	<b>Adjusted Risk Difference</b> <b>% (95% CI)</b>
<b>All-cause hospitalization ≥24 hours for acute care or death through Day 29</b>		
48 (6.8%)	68 (9.7%)	-3.0% (-5.9%, -0.1%)
<b>All-cause mortality through Day 29</b>		
1 (0.1%)	9 (1.3%)	
*The determination of primary efficacy was based on a planned interim analysis of 762 subjects. At the interim analysis, 7.3% of patients who received molnupiravir were either hospitalized or died through Day 29 (28/385), compared with 14.1% of placebo-treated patients (53/377). The adjusted risk difference was -6.8% with a 95% CI of (-11.3%, -2.4%) and 2-sided p-value = 0.0024.		
Adjusted relative risk reduction of molnupiravir compared to		


**Applicant/PHCR:** AUROGEN SA (PTY) LTD  
**Product proprietary name:** MOLNARZ  
**Dosage form and strength:** Hard capsules; 200 mg

**Submitted:** 17/05/2023

placebo for all randomized subjects was 30% (95% CI: 1%, 51%).  
 Analyses are adjusted by the stratification factor of time of COVID-19 symptom onset ( $\leq 3$  days vs.  $>3$  [4-5] days).

Efficacy results were consistent across sub-groups including age ( $> 60$  years), at risk medical conditions (e.g., obesity, diabetes), baseline COVID-19 severity (mild, moderate) and SARS-CoV-2 variants.

**Figure 1. Subgroup Efficacy Results in Non-Hospitalized Adults with COVID-19 -All- Randomized Subjects**



The corresponding confidence interval is based on Miettinen & Nurminen method. The modified intent-to-treat population is the efficacy analysis population.

**Applicant/PHCR:** AUROGEN SA (PTY) LTD

**Product proprietary name:** MOLNARZ

**Dosage form and strength:** Hard capsules; 200 mg

**Submitted:** 17/05/2023

Baseline serum samples were evaluated with the Roche Elecsys anti-N assay to test for the presence of antibodies (IgM, IgG and IgA) against the SARS-CoV-2 nucleocapsid protein.

The findings of these subgroup analyses are considered exploratory.

Higher percentages of subjects reported sustained improvement or resolution in most self-reported COVID-19 signs and symptoms, as recorded on a daily symptom diary, in the

**MOLNARZ** group compared to the placebo group.

## 5.2 Pharmacokinetic properties

Molnupiravir is a 5'-isobutyrate prodrug that is hydrolysed to NHC prior to reaching systemic circulation. The pharmacokinetics of NHC are similar in healthy subjects and patients with COVID-19.

The pharmacokinetics of NHC at steady-state following administration of 800 mg molnupiravir every 12 hours are provided below in Table 3.

**Table 3: Pharmacokinetics of NHC after administration of 800mg MOLNARZ every 12 hours**

NHC Geometric Mean (%CV)		
AUC <sub>0-12hr</sub> (ng×hr/mL) *	C <sub>max</sub> (ng/mL) †	C <sub>12hr</sub> (ng/mL) *
8260 (41.0)	2970 (16.8)	31.1 (124)
% CV: Geometric coefficient of variation. * Values were obtained from population PK analysis. † Values were obtained from a Phase 1 study of healthy subjects.		

## Absorption

Following twice daily oral administration of 800 mg molnupiravir, the median time to peak plasma NHC concentrations (T<sub>max</sub>) was 1.5 hours.

**Applicant/PHCR:** AUROGEN SA (PTY) LTD  
**Product proprietary name:** MOLNARZ  
**Dosage form and strength:** Hard capsules; 200 mg

**Submitted:** 17/05/2023

#### *Effect of Food on Oral Absorption*

In healthy subjects, the administration of a single 200 mg dose of molnupiravir with a high-fat meal had no significant effect on NHC AUC and resulted in a 35 % reduction in  $C_{max}$ .

Molnupiravir can be taken with or without food.

#### **Distribution**

NHC does not bind to plasma proteins.

#### **Elimination**

The effective half-life of NHC is approximately 3,3 hours. The fraction of dose excreted as NHC in the urine was  $\leq 3\%$  in healthy participants.

#### **Other special populations**

##### *Gender, Race, Age*

Population pharmacokinetic analysis showed that age, gender, race and ethnicity do not meaningfully influence the pharmacokinetics of NHC.

##### *Paediatric Patients*

The pharmacokinetics of molnupiravir in paediatric patients less than 18 years of age have not been evaluated.

##### *Renal Impairment*

Renal clearance is not a meaningful route of elimination for NHC. No dose adjustment in patients with any degree of renal impairment is needed. In a population PK analysis, mild or moderate renal impairment did not have a meaningful impact on the PK of NHC. While the PK of NHC has not been evaluated in patients with eGFR less than 30 mL/min/1.73 m<sup>2</sup> or on

**Applicant/PHCR:** AUROGEN SA (PTY) LTD

**Product proprietary name:** MOLNARZ

**Dosage form and strength:** Hard capsules; 200 mg

**Submitted:** 17/05/2023

dialysis, severe renal impairment and end-stage renal disease (ESRD) are not expected to have a significant effect on NHC exposure (see section 4.2 Renal Impairment).

#### *Hepatic Impairment*

The pharmacokinetics of molnupiravir and NHC has not been evaluated in patients with hepatic impairment. Preclinical data indicate that hepatic elimination is not expected to be a major route of NHC elimination therefore hepatic impairment is unlikely to affect NHC exposure. No dose adjustment in patients with hepatic impairment is needed (see section 4.2 Hepatic Impairment).

### **5.3 Preclinical safety data**

#### **General Toxicity**

Reversible, dose-related bone marrow toxicity affecting all haematopoietic cell lines was observed in dogs at  $\geq 17$  mg/kg/day (0.4 times the human NHC exposure at the recommended human dose (RHD)). Mild decreases in peripheral blood cell and platelet counts were seen after 7 days of molnupiravir treatment progressing to more severe haematological changes after 14 days of treatment. Neither bone marrow nor haematological toxicity was observed in a 1-month toxicity study in mice up to 2,000 mg/kg/day (19 times the human NHC exposure at the RHD) and a 3-month toxicity study in rats up to 1,000 mg/kg/day (9.3 and 15 times the human NHC exposure at the RHD in females and males, respectively).

Bone and cartilage toxicity, consisting of an increase in the thickness of physeal and epiphyseal growth cartilage with decreases in trabecular bone was observed in the femur and tibia of rapidly growing rats in a 3-month toxicity study at  $\geq 500$  mg/kg/day (5.4 times the human NHC exposure at the RHD). There was no bone or cartilage toxicity in a 1-month toxicity study in rapidly growing rats up to 500 mg/kg/day (4.2 and 7.8 times the human NHC

**Applicant/PHCR:** AUROGEN SA (PTY) LTD

**Product proprietary name:** MOLNARZ

**Dosage form and strength:** Hard capsules; 200 mg

**Submitted:** 17/05/2023

exposure at the RHD in females and males, respectively), in dogs dosed for 14 days up to 50 mg/kg/day (1.6 times the human NHC exposure at the RHD), or in a 1-month toxicity study in mice up to 2,000 mg/kg/day (19 times the human NHC exposure at the RHD). Growth cartilage is not present in mature skeletons; therefore the bone and cartilage findings are not relevant for adult humans. The clinical significance of these findings for paediatric patients is unknown.

### **Carcinogenesis**

Carcinogenicity studies with molnupiravir have not been conducted.

### **Mutagenesis**

Molnupiravir and NHC were positive in the *in vitro* bacterial reverse mutation assay (Ames assay) with and without metabolic activation. In 2 distinct *in vivo* rodent mutagenicity models (Pig-a mutagenicity assay and Big Blue® (cII Locus) transgenic rodent assay) molnupiravir did not induce increased mutation rates relative to untreated historical control animals, and therefore is not mutagenic *in vivo*. Molnupiravir was negative for induction of chromosomal damage in *in vitro* micronucleus (with and without metabolic activation) and *in vivo* rat micronucleus assays. Based on the totality of the genotoxicity data, molnupiravir is of low risk for genotoxicity or mutagenicity in clinical use.

### **Reproduction**

There were no effects on fertility, mating performance or early embryonic development when molnupiravir was administered to female or male rats at NHC exposures approximately 2 and 6 times, respectively, the human NHC exposure at the recommended human dose (RHD).

**Applicant/PHCR:** AUROGEN SA (PTY) LTD

**Product proprietary name:** MOLNARZ

**Dosage form and strength:** Hard capsules; 200 mg

**Submitted:** 17/05/2023

## Development

In an embryofoetal development (EFD) study in rats, molnupiravir was administered orally to pregnant rats at 0, 100, 250, or 500 mg/kg/day from gestation days (GDs) 6 to 17.

Molnupiravir was also administered orally to pregnant rats at up to 1000 mg/kg/day from GDs 6 to 17 in a preliminary EFD study. Developmental toxicities included post-implantation losses, malformations of the eye, kidney, and axial skeleton, and rib variations at 1,000 mg/kg/day (8 times the human NHC exposure at the RHD) and decreased foetal body weights and delayed ossification at  $\geq 500$  mg/kg/day (2,9 times the human NHC exposure at the RHD). There were no developmental toxicities at  $\leq 250$  mg/kg/day (0,8 times the human NHC exposure at the RHD). Maternal toxicities included decreased food consumption and body weight losses, resulting in the early sacrifice of individual animals at 1000 mg/kg/day, and decreased body weight gain at 500 mg/kg/day.

In an EFD study in rabbits, molnupiravir was administered orally to pregnant rabbits at 0, 125, 400, or 750 mg/kg/day from GDs 7 to 19. Developmental toxicity was limited to reduced foetal body weights at 750 mg/kg/day (18 times the human NHC exposures at the RHD). There was no developmental toxicity at  $\leq 400$  mg/kg/day (7 times the human NHC exposures at the RHD). Maternal toxicities included reduced food consumption and body weight gains, and abnormal faecal output at 750 mg/kg/day.

## 6. PHARMACEUTICAL PARTICULARS

### 6.1 List of excipients

The active substance is molnupiravir. The other ingredients of **MOLNARZ** are:

- Cellulose Microcrystalline PH101 Ph.Eur
- Croscarmellose Sodium Ph.Eur

**Applicant/PHCR:** AUROGEN SA (PTY) LTD  
**Product proprietary name:** MOLNARZ  
**Dosage form and strength:** Hard capsules; 200 mg

**Submitted:** 17/05/2023

- Hydroxypropyl cellulose Ph.Eur
- Magnesium stearate Ph.Eur

The capsule shell consists of the following:

- Hydroxy propylmethyl Cellulose EP
- Carrageenan EP
- Potassium Chloride USP/EP/NF
- Brilliant Blue FCF (E133)
- Iron oxide Yellow (E172)
- Titanium Dioxide Ph.Eur. (E171)

The capsule printing edible black ink consists of the following:

- Shellac (E904) Ph.Eur.
- Dehydrated Alcohol (E1510) Ph.Eur.
- Isopropyl alcohol Ph.Eur.
- Butyl Alcohol Ph.Eur.
- Propylene Glycol (E1520) Ph.Eur.
- Strong Ammonia Solution (E527) Ph.Eur.
- Potassium Hydroxide (E525) Ph.Eur.
- Purified water Ph.Eur.
- Black Iron oxide (E172) NF

## 6.2 Incompatibilities

Not applicable

**Applicant/PHCR:** AUROGEN SA (PTY) LTD  
**Product proprietary name:** MOLNARZ  
**Dosage form and strength:** Hard capsules; 200 mg

**Submitted:** 17/05/2023

### 6.3 Shelf life

Proposed: 24 months

### 6.4 Special precautions for storage

Store at or below 30 °C.

Protect from moisture.

Store in the original container. Keep tightly closed.

Keep the container in the outer carton.

### 6.5 Nature and contents of container

**MOLNARZ:**

**HDPE containers:**

**MOLNARZ** 200 mg are also proposed to be packed in white opaque round HDPE containers closed with 38 mm white opaque polypropylene continuous thread (CT) closure with induction sealing wad. This HDPE container will be further packed in preprinted carton with package leaflet.

**Pack size:** 40's

### 6.6 Special precautions for disposal of a used medicine or waste materials derived from such medicine and other handling of the product

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

**Applicant/PHCR:** AUROGEN SA (PTY) LTD  
**Product proprietary name:** MOLNARZ  
**Dosage form and strength:** Hard capsules; 200 mg

**Submitted:** 17/05/2023

## **7. HOLDER OF THE CERTIFICATE OF REGISTRATION**

AUROGEN SA (Pty) Ltd

Woodhill Office Park, Building 1, First Floor

53 Phillip Engelbrecht Avenue

Meyersdal, Ext. 12, 1448

Johannesburg

South Africa

## **8. REGISTRATION NUMBER(S)**

57/20.2.8/0169

## **9. DATE OF FIRST AUTHORISATION**

25 JULY 2023

## **10. DATE OF REVISION OF TEXT**