

**Equity Pharmaceuticals (Pty) Ltd**  
NORADRENALINE (NOREPINEPHRINE) EQUITY  
Concentrate for solution for infusion  
Registration No:57/5.1/0871  
*Each 1,0 mL solution contains noradrenaline (norepinephrine) tartrate equivalent to 2,0 mg  
noradrenaline (norepinephrine)*

**1.3.1.1.1 Approved Professional Information**  
eCTD 0007: Closing Information  
Submitted: 26 February 2026

## SCHEDULING STATUS

S4

### 1. NAME OF THE MEDICINE

NORADRENALINE (NOREPINEPHRINE) EQUITY, 1 mg/mL, concentrate for solution for infusion

### 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each mL of concentrate for solution for infusion contains 2 mg noradrenaline (norepinephrine) tartrate, equivalent to 1 mg noradrenaline (norepinephrine).

Each 4 mL ampoule contains 8 mg noradrenaline (norepinephrine) tartrate, equivalent to 4 mg noradrenaline (norepinephrine).

Each 8 mL ampoule contains 16 mg noradrenaline (norepinephrine) tartrate, equivalent to 8 mg noradrenaline (norepinephrine).

When diluted as recommended, each mL contains 80 micrograms noradrenaline (norepinephrine) tartrate, equivalent to 40 micrograms noradrenaline (norepinephrine).

#### *Excipient(s) with known effect:*

Each mL of concentrate for solution for infusion contains 3,3 mg equivalent to 0,14 mmol of sodium.

Each 4 mL ampoule contains 13,2 mg equivalent to 0,57 mmol of sodium.

Each 8 mL ampoule contains 26,4 mg equivalent to 1,14 mmol of sodium.

Sugar free.

For full list of excipients, see section 6.1.

### 3. PHARMACEUTICAL FORM

**Equity Pharmaceuticals (Pty) Ltd**  
NORADRENALINE (NOREPINEPHRINE) EQUITY  
Concentrate for solution for infusion  
Registration No:57/5.1/0871  
*Each 1,0 mL solution contains noradrenaline (norepinephrine) tartrate equivalent to 2,0 mg noradrenaline (norepinephrine)*

**1.3.1.1.1 Approved Professional Information**  
eCTD 0007: Closing Information  
Submitted: 26 February 2026

Concentrate for solution for infusion

Clear, colourless or slightly yellowish solution, practically free from visible particles.

pH = 3,0 to 4,0

Osmolality: 250 – 320 mOsm/kg

## **4. CLINICAL PARTICULARS**

### **4.1 Therapeutic indications**

NORADRENALINE (NOREPINEPHRINE) EQUITY is indicated in adults for the emergency restoration of blood pressure in cases of acute hypotension.

### **4.2 Posology and method of administration**

#### **Posology**

##### ***Adults:***

##### ***Initial rate of infusion:***

When diluted as recommended in section 6.6 (the concentration of the prepared infusion is 40 mg/litre noradrenaline (norepinephrine) (80 mg/litre noradrenaline (norepinephrine) tartrate)), the initial rate of infusion, at a body weight of 70 kg, should be between 10 mL/hour and 20 mL/hour (0,16 mL/min to 0,33 mL/min). This is equivalent to 0,4 mg/hour to 0,8 mg/hour noradrenaline (norepinephrine) (0,8 mg/hour to 1,6 mg/hour noradrenaline (norepinephrine) tartrate). Some clinicians may wish to start at a lower initial infusion rate of 5 mL/hour (0,08 mL/min), equivalent to 0,2 mg/hour noradrenaline (norepinephrine) (0,4 mg/hour noradrenaline (norepinephrine) tartrate).

##### ***Titration of dose:***

Once an infusion of NORADRENALINE (NOREPINEPHRINE) EQUITY has been established the dose should be titrated in steps of 0,05 – 0,1 µg/kg/min of noradrenaline (norepinephrine) according to the pressor effect observed. There is great individual variation in the dose required to attain and maintain normotension.

*Each 1,0 mL solution contains noradrenaline (norepinephrine) tartrate equivalent to 2,0 mg noradrenaline (norepinephrine)*

The aim should be to establish a low normal systolic blood pressure (100 – 120 mm Hg) or to achieve an adequate mean arterial blood pressure (greater than 65 – 80 mm Hg – depending on the patient’s condition).

<b>NORADRENALINE (NOREPINEPHRINE) EQUITY infusion solution 40 mg/litre (40 µg/mL)</b>			
<b>noradrenaline (norepinephrine)</b>			
<b>Patient’s weight</b>	<b>Posology (µg/kg/min)</b>	<b>Posology (mg/hour)</b>	<b>Infusion rate</b>
	<b>noradrenaline (norepinephrine)</b>	<b>noradrenaline (norepinephrine)</b>	<b>(mL/hour)</b>
50 kg	0,05	0,15	3,75
	0,1	0,3	7,5
	0,25	0,75	18,75
	0,5	1,5	37,5
	1	3	75
60 kg	0,05	0,18	4,5
	0,1	0,36	9
	0,25	0,9	22,5
	0,5	1,8	45
	1	3,6	90
70 kg	0,05	0,21	5,25
	0,1	0,42	10,5
	0,25	1,05	26,25
	0,5	2,1	52,5
	1	4,2	105
80 kg	0,05	0,24	6
	0,1	0,48	12

*Each 1,0 mL solution contains noradrenaline (norepinephrine) tartrate equivalent to 2,0 mg noradrenaline (norepinephrine)*

	0,25	1,2	30
	0,5	2,4	60
	1	4,8	120
90 kg	0,05	0,27	6,75
	0,1	0,54	13,5
	0,25	1,35	33,75
	0,5	2,7	67,5
	1	5,4	135

Some medical practitioners may prefer to dilute to other concentrations. If dilutions other than 40 mg/litre are used, check the infusion rate calculation carefully before starting treatment.

*Duration of treatment and monitoring:*

NORADRENALINE (NOREPINEPHRINE) EQUITY should be continued for as long as vasoactive medicine support is indicated. The patient should be monitored carefully for the duration of therapy. Blood pressure should be carefully monitored for the duration of therapy.

*Withdrawal of therapy:*

NORADRENALINE (NOREPINEPHRINE) EQUITY infusion should be gradually decreased since abrupt withdrawal can result in acute hypotension.

**Special populations**

*Patients with renal or hepatic impairment:*

There is no experience in treatment of renally or hepatically impaired patients.

*Elderly population:*

**Equity Pharmaceuticals (Pty) Ltd**  
NORADRENALINE (NOREPINEPHRINE) EQUITY  
Concentrate for solution for infusion  
Registration No:57/5.1/0871  
*Each 1,0 mL solution contains noradrenaline (norepinephrine) tartrate equivalent to 2,0 mg  
noradrenaline (norepinephrine)*

**1.3.1.1.1 Approved Professional Information**  
eCTD 0007: Closing Information  
Submitted: 26 February 2026

As for adults but see section 4.4.

### **Paediatric population**

The safety and efficacy of NORADRENALINE (NOREPINEPHRINE) EQUITY in children and adolescents have not been established.

### **Method of administration**

For intravenous use.

NORADRENALINE (NOREPINEPHRINE) EQUITY solution for infusion is infused as a diluted solution intravenously. To avoid ischemic necrosis (skin, extremities) a cannula placed in a sufficiently larger vein or a central venous access to the infusion should be used.

The infusion should be at a controlled rate using either a syringe pump or an infusion pump or a drip counter.

For dilution instructions see section 6.6.

### **4.3 Contraindications**

- Hypersensitivity to noradrenaline (norepinephrine) or to any of the excipients of NORADRENALINE (NOREPINEPHRINE) EQUITY as listed in section 6.1.
- Hypotension due to blood volume deficit (hypovolaemia).
- Do not use with cyclopropane and halothane anaesthetics as this may cause serious cardiac dysrhythmias including ventricular fibrillation. For interactions see section 4.5.

### **4.4 Special warnings and precautions for use**

NORADRENALINE (NOREPINEPHRINE) EQUITY should only be administered by healthcare

professionals who are familiar with its use.

***Warning:***

- Do not use undiluted.
- NORADRENALINE (NOREPINEPHRINE) EQUITY is contraindicated in hypotensive patients due to hypovolemia, however, may still be considered as a short-term emergency measure to support blood supply to coronary and cerebral arteries until general blood or solution infusion can be initiated.
- NORADRENALINE (NOREPINEPHRINE) EQUITY should be used only in conjunction with appropriate blood volume replacement.
- When infusing NORADRENALINE (NOREPINEPHRINE) EQUITY, the blood pressure and rate of flow should be checked frequently to avoid hypertension.
- The products administered by injection must always be visually inspected and cannot be used if the presence of particles or a change of colouring is noted.
- The patients receiving NORADRENALINE (NOREPINEPHRINE) EQUITY should be closely monitored in order to identify early symptoms of vasopressor-induced limb ischaemia and implement the appropriate measures (e.g. elevation of the extremity, splinting, warming of affected limb with ad hoc device, use of vasodilating agents) to prevent progression and minimize the risks associated with necrosis of the extremities.
- Extravasation risk:  

The infusion site should be checked frequently for free flow. Care should be taken to avoid extravasation that would cause a necrosis of the tissues surrounding the vein used for the injection. Because of the vasoconstriction of the vein wall with increased permeability, there might be some leakage of noradrenaline (norepinephrine) in the tissues surrounding the infused vein causing a blanching of the tissues which is not due to an obvious extravasation. Hence if blanching occurs, consideration should be given to changing the infusion site to allow the effects of local vasoconstriction to subside.

- Treatment of the ischemia due to extravasation:

During an extravascular leak of the product or an injection besides the vein, tissue destruction can appear resulting from the vasoconstrictive action of the medicine on the blood vessels. The area should be infiltrated as quickly as possible with 10 to 15 mL of physiological salt solution containing 5 to 10 mg of phentolamine mesilate, an adrenergic blocking agent. For this purpose, it is necessary to use a syringe provided with a fine needle and to inject locally throughout the area, which is easily identified by its cold, hard and pallid appearance.

***Precautions for use:***

Caution and respect of the strict indication must be retained in case of:

- Major left ventricular dysfunction associated with acute hypotension; a careful evaluation of patient's blood pressure is needed. Supportive therapy should be initiated simultaneously with diagnostic evaluation. NORADRENALINE (NOREPINEPHRINE) EQUITY should be reserved for patients with cardiogenic shock and refractory hypotension, in particular those without elevated systemic vascular resistance. It should be started at a dosage of 2 to 4 µg/min and titrated upwards and titrated as necessary. If systemic perfusion or systolic pressure cannot be maintained at > 90 mm Hg with a dosage of 15 µg/min, it is unlikely that a further increase will be beneficial.
- Particular caution should be observed in patients with coronary, mesenteric or peripheral vascular thrombosis because noradrenaline (norepinephrine) may increase the ischaemia and extend the area of infarction. Similar caution should be observed in patients with hypotension following myocardial infarction and in patients with Prinzmetal's variant angina.
- Occurrence of heart rhythm disorders during the treatment must lead to a reduction in the dosage.
- Caution is advised in patients with hyperthyroidism or diabetes mellitus.
- The elderly may be especially sensitive to the effects of noradrenaline (norepinephrine) e.g., NORADRENALINE (NOREPINEPHRINE) EQUITY, due to the greater frequency of hepatic, renal or

cardiac function and concomitant disease or other medicine therapy.

- The use of NORADRENALINE (NOREPINEPHRINE) EQUITY in children is not recommended (see section 4.2).

Perfusion of NORADRENALINE (NOREPINEPHRINE) EQUITY must be performed with continuous monitoring of blood pressure and cardiac frequency.

Prolonged administration of any potent vasopressor may result in plasma volume depletion which should be continuously corrected by appropriate fluid and electrolyte replacement therapy. If plasma volumes are not corrected, hypotension may recur when the infusion is discontinued, or blood pressure may be maintained at the risk of severe peripheral and visceral vasoconstriction (e.g., decreased renal perfusion) with diminution in blood flow and tissue perfusion with subsequent tissue hypoxia and lactic acidosis and possible ischaemic injury. Gangrene of extremities has been reported less frequently.

The vasopressor effect (resulting from the adrenergic action in the vessels) can be reduced by the concomitant administration of an alpha-blocking medicine (phentolamine mesilate) whereas the administration of a beta-blocking medicine (propranolol) may result in a reduction of the stimulating effect of the product on the heart and in an increase of the hypertensive effect (through reduction of arteriolar dilatation), resulting from beta-1-adrenergic stimulation.

In cases where it is necessary to administer NORADRENALINE (NOREPINEPHRINE) EQUITY at the same time as total blood or plasma, the latter must be administered in a separate drip.

NORADRENALINE (NOREPINEPHRINE) EQUITY contains sodium.

NORADRENALINE (NOREPINEPHRINE) EQUITY contains 3,3 mg sodium per mL, equivalent to 0,16 % of the WHO recommended maximum daily intake of 2 g sodium for an adult.

#### **4.5 Interaction with other medicines and other forms of interaction**

##### ***Contraindicated combinations***

- *Volatile halogen anaesthetics*: severe ventricular dysrhythmia (increase in cardiac excitability). The use of pressor amines with cyclopropane, halothane, chloroform, enflurane or other halogenated anaesthetics may cause serious cardiac dysrhythmias, because of the possibility of increasing the risk of ventricular fibrillation. Noradrenaline (norepinephrine) is contraindicated in combination with these medicines. See section 4.3.

##### ***Inadvisable combinations***

- *Tricyclic antidepressants*: paroxysmal hypertension with the possibility of dysrhythmia (inhibition of the entry of sympathomimetics into sympathetic fibres).
- *Serotonergic-adrenergic antidepressants*: paroxysmal hypertension with the possibility of dysrhythmia (inhibition of the entry of sympathomimetics into sympathetic fibres).
- *Desipramine*: significantly increase the toxicity of noradrenaline (norepinephrine).
- *Digitalis glycosides*: may occasionally cause dysrhythmias.
- *Levodopa*: may enhance the effects of noradrenaline (norepinephrine).
- *Antihistamines*, as some may block the intake of catecholamines by peripheral tissues and increase the toxicity of injected noradrenaline (norepinephrine).
- *Chlorpheniramine hydrochloride, tripeleminamine hydrochloride*: significantly increase the toxicity of noradrenaline (norepinephrine).

##### ***Combinations requiring precautions for use***

- *Non-selective MAO inhibitors (or within 14 days of cessation of such therapy)*: increase in the pressor action of the sympathomimetic which is usually moderate. Should only be used under close medical

supervision.

- *Selective MAO-A inhibitors*: by extrapolation from non-selective MAO inhibitors, risk of increase in the pressor action. Should only be used under close medical supervision.
- *Linezolid*: by extrapolation from non-selective MAO inhibitors, risk of increase in the pressor action. Should only be used under close medical supervision.
- *Alpha and beta blockers*: caution is required as severe hypertension may result.
- *Thyroid hormones, cardiac glycosides, anti-dysrhythmics*: caution is required as they may cause increased cardiac effects.
- *Ergot alkaloids or oxytocin*: may enhance the vasopressor and vasoconstrictive effects.
- *Desmopressin or vasopressin*: its antidiuretic effect is diminished.
- *Lithium* decreases the effect of noradrenaline (norepinephrine).
- *Guanethidine, guanadrel, reserpine, methyl dopa or amphetamine, doxapram, mazindol, rauwolfia alkaloids*: may enhance the effects of noradrenaline (norepinephrine).
- *Propofol*: concomitant administration may lead to propofol infusion syndrome (PRIS).

#### **4.6 Fertility, pregnancy and lactation**

##### **Pregnancy**

Noradrenaline (norepinephrine) may impair placental perfusion and induce foetal bradycardia. It may also exert a contractile effect on the pregnant uterus and lead to foetal asphyxia in late pregnancy.

Safety has not been established in pregnant woman.

##### **Breastfeeding**

No information is available on the use of NORADRENALINE (NOREPINEPHRINE) EQUITY in lactation. The safety of NORADRENALINE (NOREPINEPHRINE) EQUITY during breastfeeding has not been established.

**4.7 Effects on ability to drive and use machines**

Not applicable.

**4.8 Undesirable effects**

*Tabulated list of adverse reactions*

<b>System Organ Class (SOC)</b>	<b>Undesirable effect (<i>Frequency unknown</i>)</b>
<b>Metabolism and nutrition disorders</b>	Anorexia.
<b>Psychiatric disorders</b>	Anxiety, insomnia, confusion, weakness, psychotic state.
<b>Nervous system disorders</b>	Headache, tremor, lower vigilance.
<b>Eye disorders</b>	Acute glaucoma (very frequent in patients anatomically predisposed with the closing of the iridocorneal angle).
<b>Cardiac disorders</b>	Tachycardia, bradycardia (probably as a reflex result of blood pressure rising), dysrhythmias, palpitations, increase in the contractility of the cardiac muscle resulting from the beta-adrenergic effect on the heart (inotrope and chronotrope), acute cardiac insufficiency, stress cardiomyopathy.
<b>Vascular disorders</b>	Arterial hypertension and tissue hypoxia; ischemic injury due to potent vasoconstrictor action may result in coldness and paleness of the skin, members (peripheries) and the face, and gangrene of the extremities, cyanosis, hot flushes or skin redness.
<b>Respiratory, thoracic and mediastinal disorders</b>	Respiratory insufficiency or difficulty, dyspnoea.
<b>Skin and subcutaneous tissue</b>	Scarification of the skin, skin rash, hives or itching.

<b>disorders</b>	
<b>Gastrointestinal disorders</b>	Nausea, vomiting.
<b>Renal and urinary disorders</b>	Retention of urine.
<b>General disorders and administration site conditions</b>	Possibility of irritation, sloughing and necrosis at the injection site.

***Description of selected adverse reactions***

The continuous administration of vasopressor to maintain blood pressure in absence of blood volume replacement may cause the following symptoms:

- severe peripheral and visceral vasoconstriction
- decrease in renal blood flow
- decrease in urine production
- tissue hypoxia
- lactic acidosis.

In case of hypersensitivity or overdose, the following effects may appear more frequently: hypertension, photophobia, retrosternal pain, pharyngeal pain, pallor, intense sweating and vomiting.

***Reporting of suspected adverse reactions***

Reporting suspected adverse reactions after authorisation of the medicine is important. It allows continued monitoring of the benefit/risk balance of the medicine. Health care providers are requested to report any suspected adverse drug reactions to SAHPRA via the Med Safety APP (Medsafety X SAHPRA) and eReporting platform (who-umc.org) found on SAHPRA website.

**4.9 Overdose**

**Equity Pharmaceuticals (Pty) Ltd**  
NORADRENALINE (NOREPINEPHRINE) EQUITY  
Concentrate for solution for infusion  
Registration No:57/5.1/0871  
*Each 1,0 mL solution contains noradrenaline (norepinephrine) tartrate equivalent to 2,0 mg noradrenaline (norepinephrine)*

**1.3.1.1.1 Approved Professional Information**  
eCTD 0007: Closing Information  
Submitted: 26 February 2026

### Symptoms

Overdosage may result in headache, severe hypertension, reflex bradycardia, marked increase in peripheral resistance and decreased cardiac output.

These may be accompanied by violent headache, cerebral haemorrhage, photophobia, retrosternal pain, pallor, fever, intense sweating, pulmonary oedema and vomiting.

The following may also be observed: cutaneous vasoconstriction, bed sores.

### Treatment

In the event of accidental overdose, as evidenced by excessive blood pressure elevation, discontinue the medicine until the condition of the patient stabilises.

## 5. PHARMACOLOGICAL PROPERTIES

### 5.1 Pharmacodynamic properties

Category and class: A 5.1 Adrenomimetics (sympathomimetics)

Pharmacotherapeutic group: Adrenergic and Dopaminergic Agent; ATC Code: C01CA03 (C: Cardiovascular system)

#### *Mechanism of action*

Noradrenaline (norepinephrine) has a very potent action on alpha receptors and a more moderate effect on beta-1 receptors.

#### *Pharmacodynamic effects*

NORADRENALINE (NOREPINEPHRINE) EQUITY causes generalised vasoconstriction, except for the coronary vessels which it dilates indirectly by increasing the oxygen consumption. This results in an increase in the force (and in the absence of vagal inhibition) in the rate of myocardial contraction. Peripheral resistance increases, and diastolic and systolic pressures are raised.

#### *Clinical efficacy and safety*

The increase in blood pressure may cause a reflex decrease in heart rate. Vasoconstriction may result in decreased blood flow in kidneys, liver, skin and smooth muscles. Local vasoconstriction may cause haemostasis and/or necrosis.

The effect on blood pressure disappears 1 – 2 minutes after stopping the infusion.

## **5.2 Pharmacokinetic properties**

Two stereoisomers of noradrenaline (norepinephrine) exist, the biologically active L-isomer is the one present in NORADRENALINE (NOREPINEPHRINE) EQUITY.

### **Absorption**

- After intravenous administration noradrenaline (norepinephrine) has a plasmatic half-life of about 1 to 2 minutes.

### **Distribution**

- Noradrenaline (norepinephrine) is rapidly cleared from plasma by a combination of cellular reuptake and metabolism. It does not readily cross the blood-brain barrier.

### **Biotransformation**

- Methylation by catechol-o-methyltransferase
- Deamination by monoamine oxidase (MAO)
- Ultimate metabolites from both is 4-hydroxy-3-methoxymandelic acid
- Intermediate metabolites include normetanephrine and 3,4- dihydroxymandelic acid.

## **Elimination**

- Noradrenaline (norepinephrine) is mainly eliminated as glucuronide or sulphate conjugates of the metabolites in the urine.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Sodium chloride

Hydrochloric acid (for pH adjustment) or

Sodium hydroxide (for pH adjustment)

Water for injections

### **6.2 Incompatibilities**

NORADRENALINE (NOREPINEPHRINE) EQUITY must not be mixed with other medicines except those mentioned in the section 6.6.

### **6.3 Shelf life**

24 months

*After dilution:*

Chemical and physical in-use stability of diluted product (in 5 % dextrose, sodium chloride 9 mg/mL (0,9 %), or isotonic dextrose saline) has been demonstrated for 48 hours at 30 °C.

However, from a microbiological point of view, the diluted product should be used immediately. If not used immediately, in-use storage times and conditions prior to use are the responsibility of the user and would normally not be longer than 24 hours at 2 °C to 8 °C, unless dilution has taken place in controlled and validated aseptic conditions.

**Equity Pharmaceuticals (Pty) Ltd**  
NORADRENALINE (NOREPINEPHRINE) EQUITY  
Concentrate for solution for infusion  
Registration No:57/5.1/0871  
*Each 1,0 mL solution contains noradrenaline (norepinephrine) tartrate equivalent to 2,0 mg noradrenaline (norepinephrine)*

**1.3.1.1.1 Approved Professional Information**  
eCTD 0007: Closing Information  
Submitted: 26 February 2026

#### **6.4 Special precautions for storage**

Store at or below 25 °C.

Store in the original package to protect from light.

For storage conditions after dilution of the medicinal product, see section 6.3.

#### **6.5 Nature and contents of container**

NORADRENALINE (NOREPINEPHRINE) EQUITY is packaged in type I clear glass, self-breaking (one point cut) ampoules of 5 mL and 10 mL, filled to 4 mL and 8 mL, respectively.

The glass ampoules are packed into carton boxes containing 10, 50 or 100 ampoules.

Not all pack sizes may be marketed.

#### **6.6 Special precautions for disposal and other handling**

For single use.

NORADRENALINE (NOREPINEPHRINE) EQUITY should be diluted prior to intravenous infusion, either with dextrose 5%, sodium chloride 9 mg/mL (0.9 %), or with isotonic dextrose saline. It should not be mixed with other medicines.

##### *Dilution instructions:*

Add 2 mL of NORADRENALINE (NOREPINEPHRINE) EQUITY to 48 mL 5 % dextrose (or sodium chloride 9 mg/mL (0,9 %), or isotonic dextrose saline) for administration by syringe pump, or add 20 mL of NORADRENALINE (NOREPINEPHRINE) EQUITY to 480 mL 5 % dextrose (or sodium chloride 9 mg/mL (0,9 %), or isotonic dextrose saline) for administration by drip counter.

In both the cases, the final concentration of the infusion solution is usually 40 mg/litre noradrenaline (norepinephrine) (80 mg/litre noradrenaline (norepinephrine) tartrate). If other dilutions are used check the calculation carefully before starting treatment.

**Equity Pharmaceuticals (Pty) Ltd**  
NORADRENALINE (NOREPINEPHRINE) EQUITY  
Concentrate for solution for infusion  
Registration No:57/5.1/0871  
*Each 1,0 mL solution contains noradrenaline (norepinephrine) tartrate equivalent to 2,0 mg  
noradrenaline (norepinephrine)*

**1.3.1.1.1 Approved Professional Information**  
eCTD 0007: Closing Information  
Submitted: 26 February 2026

NORADRENALINE (NOREPINEPHRINE) EQUITY is compatible with PVC infusion bags.

Do not use an opened ampoule.

This product should be visually inspected prior to administration. Only a clear, colourless or slightly yellowish solution, free of particles or precipitates should be used. The ampoules with a pink colour or darker than pale yellow, or containing a precipitate should not be administered.

Any unused product or waste material should be disposed of in accordance with local requirements.

## **7. HOLDER OF CERTIFICATE OF REGISTRATION**

Equity Pharmaceuticals (Pty) Ltd

100 Sovereign Drive

Route 21 Corporate Park

Nellmapius Drive

Irene, Pretoria

Tel: +27 (0)12 345 1747

## **8. REGISTRATION NUMBER(S)**

57/5.1/0871

## **9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

20 February 2024

## **10. DATE OF REVISION OF THE TEXT**

23 February 2026