

APPROVED PROFESSIONAL INFORMATION

SCHEDULING STATUS

S3

1. NAME OF THE MEDICINE

RASSETOM 250 mg film coated tablets

RASSETOM 500 mg film coated tablets

RASSETOM 750 mg film coated tablets

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each RASSETOM 250 mg tablet contains 250 mg levetiracetam.

Each RASSETOM 500 mg tablet contains 500 mg levetiracetam.

Each RASSETOM 750 mg tablet contains 750 mg levetiracetam.

RASSETOM tablets are sugar free.

For the full list of excipients, see section 6.1.

3. PHARMACEUTICAL FORM

Film coated tablet.

RASSETOM 250 mg:

Blue, oblong-shaped, biconvex film coated tablets debossed with '250'
on one side and a score line on the other side.

RASSETOM 500 mg:

Yellow, oblong-shaped, biconvex film coated tablets debossed with

APPROVED PROFESSIONAL INFORMATION

'500' on one side and a score line on the other side.

RASSETOM 750 mg:

Peach coloured, oblong-shaped, biconvex film coated tablets

debossed with '750' on one side and a score line on the other side.

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

RASSETOM is indicated as monotherapy in the treatment of partial onset seizures with or without secondary generalisation in patients from 16 years of age with newly diagnosed epilepsy.

RASSETOM is indicated as adjunctive therapy:

- in the treatment of partial onset seizures with or without secondary generalisation in adults and children over 16 years of age with epilepsy
- in the treatment of myoclonic seizures in adults and adolescents from 12 years of age with Juvenile Myoclonic Epilepsy
- in the treatment of primary generalised tonic-clonic seizures in adults and children from 16 years of age with idiopathic generalised epilepsy.

4.2 Posology and method of administration

Posology

APPROVED PROFESSIONAL INFORMATION

Monotherapy:

Adults and adolescents from 16 years of age:

The starting dose is 250 mg twice daily, which should be increased to an initial therapeutic dose of 500 mg twice daily after two weeks. The dose can be further increased by 250 mg twice daily, every two weeks, depending on the clinical response. The maximum daily dose is 1 500 mg twice daily.

Add-on therapy:

Adults and adolescents older than 12 years (weighing 50 kg or more when indicated (see section 4.1)):

As adjunctive therapy, the therapeutic dose is 500 mg twice daily. The dose can be started on the first day of treatment. Depending upon the clinical response and tolerance, the daily dose can be increased up to 1 500 mg twice daily. Dose changes can be made in 500 mg twice daily increments or decrements every two to four weeks. The maximum daily dose is 3 000 mg.

Special populations

Elderly:

Adjustment of the dose is recommended in elderly patients with compromised renal function (see *Patients with renal impairment* below).

APPROVED PROFESSIONAL INFORMATION

Patients with renal impairment:

The RASSETOM dose should be individualised according to renal function. Refer to the table below and adjust the dose as indicated.

Dosing adjustment for adult patients with impaired renal function:

| Group | Creatinine clearance (mL/min) | Dosage and frequency |
|---|--------------------------------------|---|
| Normal | > 80 | 500 to 1 500 mg twice daily |
| Mild | 50 – 79 | 500 to 1 000 mg twice daily |
| Moderate | 30 – 49 | 250 to 750 mg twice daily |
| Severe | < 30 | 250 to 500 mg twice daily |
| End-stage renal disease patients undergoing dialysis ¹ | — | 500 to 1 000 mg once daily ² |

¹ A 750 mg loading dose is recommended on the first day of treatment with RASSETOM.

² Following dialysis, a 250 mg to 500 mg supplemental dose is recommended.

To use this dosing table, an estimate of the patient's creatinine clearance (eCL_{cr}) in mL/min is needed.

APPROVED PROFESSIONAL INFORMATION

The CL_{cr} can be estimated from the serum creatinine (S_{cr}) concentration using the modified formula of Cockfort and Gault (for use in adults):

$$\frac{eCL_{cr} \text{ (mL/min)} = [140 - \text{age}] \times \text{Wt (kg)} \times \text{constant}^*}{S_{cr} \text{ (}\mu\text{mol/L)}}$$

*Constant = 1,23 for males and 1,04 for females (0,85 x 1,23 = 1,04)

Patients with hepatic impairment:

No dose adjustment is needed in patients with mild to moderate hepatic impairment. In patients with severe hepatic impairment, the creatinine clearance may underestimate the renal insufficiency. Therefore a 50 % reduction of the daily maintenance dose is recommended when the creatinine clearance is < 70 mL/min.

Paediatric population

Monotherapy

Infants and children under the age of 12 years:

The safety and efficacy of RASSETOM in children and adolescents below 16 years as monotherapy treatment have not been established.

No data are available.

Add-on therapy

Infants and children under the age of 12 years:

APPROVED PROFESSIONAL INFORMATION

There are insufficient data available for the use of RASSETOM in children under the age of 12 years.

Adolescents (12 to 17 years) weighing less than 50 kg, when indicated (see section 4.1):

The initial therapeutic dose is 10 mg/kg twice daily. This dose can be started on the first day of treatment. Depending upon the clinical response and tolerability, the dose can be increased up to 30 mg/kg twice daily. Dose changes should not exceed increases or decreases of 10 mg/kg twice daily every two weeks. The lowest effective dose should be used. Dosage in children 50 kg or greater is the same as in adults. The doctor should prescribe the most appropriate pharmaceutical form and strength according to weight and dose.

Recommended dosage for children and adolescents with normal renal function:

| Weight | Starting dose (10 mg/kg twice daily) | Maximum dose (30 mg/kg twice daily) |
|---------------|---|--|
| 15 kg* | 150 mg twice daily | 450 mg twice daily |
| 20 kg* | 200 mg twice daily | 600 mg twice daily |
| 25 kg | 250 mg twice daily | 750 mg twice daily |
| From 50 kg** | 500 mg twice daily | 1 500 mg twice daily |

* Another formulation of levetiracetam should be used.

APPROVED PROFESSIONAL INFORMATION

** Dosage in children and adolescents 50 kg or more is the same as in adults.

Method of administration

The film coated tablets should be taken orally, swallowed with liquid, and may be taken with or without food. The daily dose is administered in two equally divided doses.

Missed dose

If a dose is missed, the tablet should be taken as soon as the missed dose is remembered. Two tablets should not be taken to make up for the missed dose.

4.3 Contraindications

- hypersensitivity to levetiracetam or other pyrrolidine derivatives or to any of the ingredients of RASSETOM
- pregnancy and lactation.

4.4 Special warnings and precautions for use

Suicide

Suicide, suicide attempt, suicidal ideation and behaviour have been reported in patients treated with anti-epileptic medicines (including levetiracetam as in RASSETOM). Analysis of randomized placebo-controlled trials of anti-epileptic medicines has shown a small

APPROVED PROFESSIONAL INFORMATION

increased risk of suicidal thoughts and behaviour. The mechanism of this risk is not known. Patients should be monitored for signs of depression and/or suicidal ideation and behaviours and appropriate treatment should be considered. Patients (and caregivers of patients) should be advised to seek medical advice should signs of depression and/or suicidal ideation or behaviour emerge (see section 4.8).

Renal impairment

Reduced doses of RASSETOM are recommended for patients with renal impairment (see section 4.2).

Patients receiving dialysis may be given a loading dose of 750 mg when starting RASSETOM followed by doses of 500 to 1 000 mg once daily, a supplemental dose of 250 to 500 mg is recommended after dialysis. In patients with severely impaired hepatic function, assessment of renal function is recommended before dose selection (see section 4.2).

Acute kidney injury

The use of RASSETOM has been very rarely associated with acute kidney injury, with a time to onset ranging from a few days to several months.

Severe hepatic impairment

RASSETOM should be used with caution in patients with severe

APPROVED PROFESSIONAL INFORMATION

hepatic impairment. No dose adjustment is needed in patients with mild to moderate hepatic impairment.

In patients with severe hepatic impairment, see section 4.2.

Blood cell counts

Rare cases of decreased blood cell counts (neutropenia, agranulocytosis, leukopenia, thrombocytopenia, and pancytopenia) have been described in association with RASSETOM administration, generally at the beginning of the treatment. Complete blood cell counts are advised in patients experiencing important weakness, pyrexia, recurrent infections, or coagulation disorders (see section 4.8).

Drug reaction with eosinophilia and systemic symptoms

(DRESS)

Drug reactions with eosinophilia and systemic symptoms (DRESS), also known as hypersensitivity syndrome (HSS), is associated with a variable pattern of systemic symptoms including fever, lymphadenopathy, facial oedema, abnormalities of the blood, liver and kidney. Other common manifestations include arthralgias, jaundice, hepatomegaly, leucocytosis and eosinophilia. It is important to note that early manifestations of hypersensitivity, such as fever and lymphadenopathy, may be present even though rash is not evident. Patients at higher risk of developing HSS/DRESS include black

APPROVED PROFESSIONAL INFORMATION

patients, patients who have experience the syndrome in the past, patients who have a family history of this syndrome and immune-suppressed patients. This syndrome is more severe in previously sensitized individuals.

If any signs and symptoms are present, the patient should be evaluated immediately, and treatment discontinued if an alternate treatment cannot be established.

Abnormal and aggressive behaviours

RASSETOM may cause psychotic symptoms and behavioural abnormalities including irritability and aggressiveness. Patients treated with RASSETOM should be monitored for developing psychiatric signs suggesting important mood and/or personality changes. If such behaviours are noticed, treatment adaptation or gradual discontinuation should be considered. If discontinuation is considered, please refer to section 4.2.

Withdrawal of treatment

Withdrawal of RASSETOM or transition to or from another type of anti-epileptic therapy should be made gradually (e.g., 500 mg twice daily decrements every two to four weeks in adults; 10 mg/kg twice daily decrements every two weeks in children) to avoid precipitating an increase in the frequency of seizures.

APPROVED PROFESSIONAL INFORMATION

Worsening of seizures

As with other types of antiepileptic drugs, RASSETOM may rarely exacerbate seizure frequency or severity. This paradoxical effect was mostly reported within the first month after levetiracetam initiation or increase of the dose, and was reversible upon drug discontinuation or dose decrease. Patients should be advised to consult their doctor immediately in case of aggravation of epilepsy.

Lack of efficacy or seizure worsening has for example been reported in patients with epilepsy associated with sodium voltage-gated channel alpha subunit 8 (SCN8A) mutations.

Electrocardiogram QT interval prolongation

Rare cases of ECG QT interval prolongation have been observed during the post-marketing surveillance. RASSETOM should be used with caution in patients with QTc-interval prolongation, in patients concomitantly treated with drugs affecting the QTc-interval, or in patients with relevant pre-existing cardiac disease or electrolyte disturbances.

Paediatric population

Available data in children did not suggest impact on growth and puberty. However, long term effects on learning, intelligence, growth, endocrine function, puberty, and childbearing potential in children remain unknown.

APPROVED PROFESSIONAL INFORMATION

4.5 Interaction with other medicines and other forms of interaction

- potential interactions with existing anti-epileptic medicines (such as phenytoin, carbamazepine, valproic acid, phenobarbital, lamotrigine, gabapentin, and primidone) have not been demonstrated in adults. As in adults, there is no evidence of clinically significant medicinal product interactions in paediatric patients receiving up to 60 mg/kg/day levetiracetam.
- a retrospective assessment of pharmacokinetic interactions in children and adolescents with epilepsy (4 to 17 years) confirmed that adjunctive therapy with orally administered levetiracetam did not influence the steady-state serum concentrations of concomitantly administered carbamazepine and valproate. However, data suggested a 20 % higher levetiracetam clearance in children taking enzyme-inducing antiepileptic medicines. Dose adjustment is not required.
- probenecid (500 mg daily), a renal tubular secretion blocking agent, has been shown to inhibit the renal clearance of the primary metabolite but not of levetiracetam. Nevertheless, the concentration of this metabolite remains low. It is expected that other medicines excreted by active tubular secretion could also reduce the renal clearance of the metabolite. The effect of

APPROVED PROFESSIONAL INFORMATION

levetiracetam on probenecid was not studied. The effect of levetiracetam on other actively secreted medicines e.g., NSAIDs and sulphonamides is unknown.

- concomitant administration of levetiracetam and methotrexate has been reported to decrease methotrexate clearance, resulting in increased/prolonged blood methotrexate concentration to potentially toxic levels. Blood methotrexate and levetiracetam levels should be carefully monitored in patients treated concomitantly with the two medicines.
- RASSETOM did not influence the pharmacokinetics of oral contraceptives (ethinyl oestradiol and levonorgestrel), endocrine parameters [luteinising hormone (LH) and progesterone] were not modified. RASSETOM 2 000 mg daily did not influence the pharmacokinetics of digoxin and warfarin, prothrombin times were not modified. Co-administration with digoxin, oral contraceptives or warfarin had no influence on the pharmacokinetics of levetiracetam.
- no information on the influence of antacids on the absorption of levetiracetam is available.
- there have been isolated reports of decreased levetiracetam efficacy when the osmotic laxative macrogol has been concomitantly administered with oral levetiracetam. Therefore, macrogol should not be taken orally for one hour before and for one hour after taking levetiracetam.

APPROVED PROFESSIONAL INFORMATION

- the extent of absorption of levetiracetam was not altered by food, but the rate of absorption was slightly reduced.
- no information on the interaction of levetiracetam with alcohol is available.

4.6 Fertility, pregnancy and lactation

Women of childbearing potential

Specialist advice should be given to women who are of childbearing potential. Treatment with levetiracetam should be reviewed when a woman is planning to become pregnant. As with all antiepileptic medicines, sudden discontinuation of levetiracetam should be avoided as this may lead to breakthrough seizures that could have serious consequences for the woman and the unborn child. Monotherapy should be preferred whenever possible because therapy with multiple antiepileptic medicines AEDs could be associated with a higher risk of congenital malformations than monotherapy, depending on the associated antiepileptics.

Pregnancy

RASSETOM is contraindicated in pregnancy and lactation (see section 4.3).

Breastfeeding

Levetiracetam is excreted in human breast milk. Patients using

APPROVED PROFESSIONAL INFORMATION

RASSETOM must not breastfeed.

Fertility

No impact on fertility was detected in animal studies (see section 5.3).

No clinical data are available and potential risk for humans is unknown.

4.7 Effects on ability to drive and use machines

RASSETOM has minor or moderate influence on the ability to drive and use machines.

Caution is recommended in patients performing skilled tasks, e.g. driving vehicles or operating machinery. At the beginning of treatment or after a dosage increase, some patients may experience somnolence or other CNS related symptoms such as headache, fatigue and dizziness.

4.8 Undesirable effects

a. Summary of the safety profile

The most frequently reported adverse reactions were nasopharyngitis, somnolence, headache, fatigue, and dizziness. The safety profile of levetiracetam is generally similar across age groups (adult and paediatric patients) and across the approved epilepsy indications.

b. Tabulated list of adverse effects

APPROVED PROFESSIONAL INFORMATION

| System Organ Class | Frequency | Side effects |
|---|-------------------------------|--|
| Infections and infestations | Frequent Less frequent | Nasopharyngitis Infection |
| Blood and lymphatic system disorders | Less frequent | Leukopenia*, neutropenia*, pancytopenia*, thrombocytopenia*, agranulocytosis |
| Immune system disorders | Less frequent | Drug reaction with eosinophilia and systemic symptoms (DRESS), Hypersensitivity (including angioedema and anaphylaxis) |
| Metabolism and nutrition disorders | Frequent Less frequent | Anorexia Decreased weight*, increased weight, hyponatraemia |
| Psychiatric disorders | Frequent Less frequent | Anxiety, depression, hostility, aggression, insomnia, nervousness, irritability Psychotic disorders*, mood swings*, agitation, personality disorder, emotional lability, suicide attempt*, suicidal ideation*, completed suicide*, abnormal behaviour*, hallucination*, anger*, confusional state*, panic attack affect/liability mood swings, thinking abnormal*, delirium, obsessive compulsive disorder (OCD)* |

APPROVED PROFESSIONAL INFORMATION

| | | |
|--|---------------|--|
| Nervous system disorders | Frequent | Dizziness, headache, somnolence, convulsion, balance disorder, lethargy, tremor |
| | Less frequent | Paraesthesia*, amnesia, memory impairment, abnormal coordination, ataxia, disturbance in attention, choreoathetosis, dyskinesia, hyperkinesia, gait disturbance, encephalopathy, seizures aggravated |
| Eye disorders | Frequent | Diplopia, blurred vision |
| Ear and labyrinth disorders | Frequent | Vertigo |
| Cardiac disorders | Less frequent | Electrocardiogram QT prolonged |
| Respiratory, thoracic and mediastinal disorders | Frequent | Pharyngitis, rhinitis, cough |
| | Less frequent | Sinusitis |
| Gastrointestinal disorders | Frequent | Diarrhoea, dyspepsia, nausea, vomiting, abdominal pain |
| | Less frequent | Pancreatitis |
| Hepatobiliary disorders | Less frequent | Abnormal liver function tests*, hepatic failure*, hepatitis* |
| Skin and subcutaneous tissue disorders | Frequent | Rash |
| | Less frequent | Alopecia, eczema*, pruritus*, toxic epidermal necrolysis*, Stevens-Johnson syndrome*, erythema multiforme* |
| Musculoskeletal, connective tissue and bone disorders | Frequent | Ataxia, muscular weakness, myalgia |
| | Less frequent | Rhabdomyolysis and blood creatine phosphokinase increased** |

APPROVED PROFESSIONAL INFORMATION

| | | |
|---|---------------|---------------------|
| Renal and urinary disorders | Less frequent | Acute kidney injury |
| General disorders and administrative site conditions | Frequent | Asthenia, fatigue |
| Investigations | Less frequent | Injury |

*Post marketing side effects

** Prevalence is significantly higher in Japanese patients when compared to non-Japanese patients

c. Description of selected adverse reactions

The risk of anorexia is higher when levetiracetam is co-administered with topiramate.

In several cases of alopecia, recovery was observed when levetiracetam was discontinued.

Bone marrow suppression was identified in some of the cases of pancytopenia.

Cases of encephalopathy generally occurred at the beginning of the treatment (a few days to a few months) and were reversible after treatment discontinuation.

d. Paediatric population

The adverse reaction profile of levetiracetam is generally similar across age groups and across the approved epilepsy indications. Safety results in paediatric patients in placebo-controlled clinical studies were consistent with the safety profile of levetiracetam in

APPROVED PROFESSIONAL INFORMATION

adults except for behavioural and psychiatric adverse reactions which were more common in children than in adults.

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicine is important. It allows continued monitoring of the benefit/risk balance of the medicine. Healthcare professionals are requested to report any suspected adverse drug reactions to SAHPRA via the Med Safety APP (Medsafety X SAHPRA) and eReporting platform (who-umc.org) found on SAHPRA website.

An email can be sent directly to the company, pharmacovigilance@pharmadynamics.co.za, to ensure safety of the product.

Signs and symptoms

Experience with levetiracetam overdose is limited. Symptoms that may occur with RASSETOM overdose include drowsiness, aggression, agitation, coma, depressed level of consciousness, respiratory depression, and somnolence.

Management of overdose

There is no specific antidote for RASSETOM overdose. Treatment should be symptomatic and supportive. Emesis should be attempted if indicated.

APPROVED PROFESSIONAL INFORMATION

Standard haemodialysis should be considered, particularly, in selected patients based on clinical state of renal impairment.

The dialyser extraction efficiency is 60 % for levetiracetam and 74 % for the primary metabolite.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Antiepileptics, other antiepileptics.

ATC code: N03AX14

Pharmacological classification: A 2.5 Anticonvulsants, including anti-epileptics.

Mechanism of action

Levetiracetam is an anti-epileptic medicine. It is a pyrrolidine derivative, the S-enantiomer of α -ethyl-2-oxo-1-pyrrolidine acetamide, which is chemically unrelated to existing anti-epileptic active substances. Levetiracetam inhibits partial and secondarily generalised tonic-clonic seizures in the kindling model of epilepsy. It is ineffective against maximum electroshock and pentylenetetrazol induced seizures. It is found to be more clinically effective against partial and secondarily tonic-clonic seizures. The mechanism by which levetiracetam exerts these anti-seizure effects is unknown. No evidence for an action on voltage-gated sodium Na⁺ channels or either gamma-aminobutyric acid GABA- or glutamate-mediated

APPROVED PROFESSIONAL INFORMATION

synaptic transmission has emerged.

5.2 Pharmacokinetic properties

The pharmacokinetic profile is dose linear with low intra- and inter-subject variability. There is no modification of the clearance after repeated administration. There is no evidence for any relevant gender, race, or circadian variability. The pharmacokinetic profile is comparable in healthy volunteers and in patients with epilepsy.

Due to its complete and linear absorption, plasma levels can be predicted from the oral dose of levetiracetam expressed as mg/kg bodyweight. Therefore, there is no need for plasma level monitoring of levetiracetam.

A significant correlation between saliva and plasma concentrations has been shown in adults and children (ratio of saliva/plasma concentrations ranged from 1 to 1,7 for oral tablet formulation and after 4 hours post-dose for oral solution formulation).

Absorption:

Levetiracetam is rapidly and almost completely absorbed after oral administration, oral bioavailability is 100 %. Peak plasma concentrations (C_{max}) are achieved at 1,3 hours after dosing. Steady-state is achieved after 2 days of a twice daily administration schedule.

Plasma protein binding is minimal (< 10 %). Peak concentrations

APPROVED PROFESSIONAL INFORMATION

(C_{max}) are typically 31 and 43 µg/mL following a single 1 000 mg dose and repeated 1 000 mg twice daily dose, respectively. The extent of absorption is dose-independent and is not altered by food.

Distribution:

No tissue distribution data are available in humans. Neither levetiracetam nor its primary metabolite are significantly bound to plasma proteins (< 10 %). The volume of distribution of levetiracetam is approximately 0,5 to 0,7 L/kg, a value close to the volume of distribution of intracellular and extracellular water.

Biotransformation:

Levetiracetam is not extensively metabolised in humans.

The major metabolic pathway (24 % of the dose) is an enzymatic hydrolysis of the acetamide group. Production of the primary metabolite, ucb L057, is not supported by liver cytochrome P450 isoforms. Hydrolysis of the acetamide group was measurable in a large number of tissues including blood cells. The metabolite ucb L057 is pharmacologically inactive.

Two minor metabolites were also identified. One was obtained by hydroxylation of the pyrrolidone ring (1.6 % of the dose) and the other one by opening of the pyrrolidone ring (0.9 % of the dose).

Other unidentified components accounted only for 0.6 % of the dose.

APPROVED PROFESSIONAL INFORMATION

No enantiomeric interconversion was evidenced *in vivo* for either levetiracetam or its primary metabolite.

In vitro, levetiracetam and its primary metabolite have been shown not to inhibit the major human liver cytochrome P₄₅₀ isoforms (CYP3A4, 2A6, 2C9, 2C19, 2D6, 2E1 and 1A2), glucuronyl transferase (UGT1A1 and UGT1A6) and epoxide hydroxylase activities. In addition, levetiracetam does not affect the *in vitro* glucuronidation of valproic acid.

In human hepatocytes in culture, levetiracetam had little or no effect on CYP1A2, SULT1E1 or UGT1A1. Levetiracetam caused mild induction of CYP2B6 and CYP3A4. The *in vitro* data and *in vivo* interaction data on oral contraceptives, digoxin and warfarin indicate that no significant enzyme induction is expected *in vivo*. Therefore, the interaction of levetiracetam with other substances, or vice versa, is unlikely.

Elimination:

The plasma half-life in adults has been reported as 7 ± 1 hours and did not vary with dose, route of administration or repeated administration. The total body clearance was a mean of 0, 96 mL/min/kg. Approximately 95 % are excreted in the urine, (approximately 93 % of the dose was excreted within 48 hours) 66 %

APPROVED PROFESSIONAL INFORMATION

of which is unchanged and 24 % is metabolised by hydrolysis of the acetamide group.

Excretion via faeces accounted for only 0.3 % of the dose.

Levetiracetam neither induces nor is a high-affinity substrate for cytochrome P450 isoforms of glucuronidation enzymes and thus is devoid of known interactions with other anti-seizure medicines, oral contraceptives, or anticoagulants.

Pharmacokinetics in special patient groups

Elderly:

In the elderly, the half-life is increased by about 40 % (10 to 11 hours). This is related to the decrease in renal function in this population (see section 4.2).

Renal impairment:

The apparent body clearance of both levetiracetam and of its primary metabolite is correlated to the creatinine clearance. It is therefore recommended that the maintenance daily dose of levetiracetam be adjusted, based on creatinine clearance in patients with moderate and severe renal impairment (see section 4.2). In anuric end-stage renal disease subjects, the half-life was approximately 25 and 3,1 hours during inter-dialytic and intradialytic periods, respectively. The fractional removal of levetiracetam was 51 % during a typical 4-hour dialysis session.

Hepatic impairment:

APPROVED PROFESSIONAL INFORMATION

In subjects with mild and moderate hepatic impairment, there was no relevant modification of the clearance of levetiracetam. In most subjects with severe hepatic impairment, the clearance of levetiracetam was reduced by more than 50 % due to a concomitant renal impairment (see section 4.2).

5.3 Preclinical safety data

Non-clinical data reveal no special hazard for humans based on conventional studies of safety pharmacology, genotoxicity, and carcinogenic potential.

Adverse effects, not observed in clinical studies but seen in the rat and to a lesser extent in the mouse, at exposure levels similar to human exposure levels and with possible relevance for clinical use were liver changes, indicating an adaptive response such as increased weight and centrilobular hypertrophy, fatty infiltration, and increased liver enzymes in plasma.

No adverse reactions on male or female fertility or reproduction performance were observed in rats at doses up to 1800 mg/kg/day (x 6 the MRHD on a mg/m² or exposure basis) in parents and F1 generation.

APPROVED PROFESSIONAL INFORMATION

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Tablet cores:

Croscarmellose sodium

Magnesium stearate

Maize starch

Microcrystalline cellulose

Povidone

Silica colloidal

Talc

Tablet coating:

Opadry blue AMB 84F80803: (FD & C blue #2/Indigo carmine
aluminium lake)

Opadry pink AMB 84F 84674: (FC & C yellow #6 Sunset yellow FCF
aluminium lake, iron oxide red, iron oxide yellow)

Opadry white AMB 84F58775

Opadry yellow AMB 84F82508: (Iron oxide yellow)

Excipients common to Opadry variants:

Macrogol 3350

Macrogol 6000

Polyvinyl alcohol (part hydrolysed)

Talc

Titanium dioxide

APPROVED PROFESSIONAL INFORMATION

6.2 Incompatibilities

Not applicable.

6.3 Shelf life

3 years.

6.4 Special precautions for storage

Store at or below 30 °C. Keep blisters in carton until required for use.

6.5 Nature and contents of container

Clear PVC/ Aluminium blister strips containing 10 tablets. Six blister strips (6 x 10) or three blister strips (3 x 10) are packed in an outer cardboard box.

6.6 Special precautions for disposal

No special requirements.

7. HOLDER OF THE CERTIFICATE OF REGISTRATION

Pharma Dynamics (Pty) Ltd

1st Floor, Grapevine House, Steenberg Office Park

Silverwood Close

Westlake, Cape Town

7945, South Africa

Tel.: +27 21 707 7000

APPROVED PROFESSIONAL INFORMATION

or 0860-PHARMA (742 762)

8. REGISTRATION NUMBER(S)

RASSETOM 250 mg: A44/2.5/0368

RASSETOM 500 mg: A44/2.5/0369

RASSETOM 750 mg: A44/2.5/0370

9. DATE OF FIRST AUTHORISATION

Date of registration: 01 March 2013

10. DATE OF REVISION OF THE TEXT

28 February 2025

NAMIBIA:

DYNA LEVETIRACETAM 250 mg: NAM NS2 13/2.5/0178

DYNA LEVETIRACETAM 500 mg: NAM NS2 13/2.5/0179

DYNA LEVETIRACETAM 750 mg: NAM NS2 13/2.5/0180