

**Applicant/PHCR:** AUROGEN SOUTH AFRICA (PTY) LTD  
**Product proprietary name:** SHILOVA 500 mg/ 1000 mg TABLETS  
**Dosage form and strength:** TABLETS 500 mg/ 1000 mg

**Date:** 13/02/2025

## PROFESSIONAL INFORMATION

**SCHEDULING STATUS:** S4

### 1 NAME OF THE MEDICINE

**SHILOVA 500 mg TABLETS** (tablets)

**SHILOVA 1 g TABLETS** (tablets)

### 2 QUALITATIVE AND QUANTITATIVE COMPOSITION

#### **SHILOVA 500 mg TABLETS:**

Each film-coated tablet contains valaciclovir hydrochloride equivalent to 500 mg valaciclovir.

#### **SHILOVA 1 g TABLETS:**

Each film-coated tablet contains valaciclovir hydrochloride equivalent to 1 g valaciclovir.

For full list of excipients, see section 6.1.

Sugar free

### 3 PHARMACEUTICAL FORM

#### **SHILOVA 500 mg TABLETS:**

Blue, film coated, capsule shaped tablets with a partial score bar on both sides containing "F" on one side and "9" and "3" on the other side.

#### **SHILOVA 1 g TABLETS:**

Blue, film coated, capsule shaped tablets with a partial score bar on both sides containing "F" on one side and "8" and "3" on the other side.

## 4 CLINICAL PARTICULARS

### 4.1 Therapeutic indication

**SHILOVA TABLETS** is indicated for the treatment of herpes zoster (shingles). **SHILOVA TABLETS** reduces the duration of zoster-associated pain, which includes acute and postherpetic neuralgia, thus accelerating resolution of pain. **SHILOVA TABLETS** also reduces the proportion of patients with zoster-associated pain.

**SHILOVA TABLETS** is indicated for the episodic treatment of recurrent genital herpes in immunocompetent adult patients.

**SHILOVA TABLETS** is indicated for the prevention (suppression) of recurrent herpes simplex infection of the skin and mucous membrane of the ano-genital area.

**SHILOVA TABLETS** is indicated for the prophylaxis of cytomegalovirus (CMV) infection, CMV disease and other herpes virus infections following organ transplantation, where special risk exists.

### 4.2 Posology and method of administration

#### Posology

##### Dosage in adults:

*For the treatment of Herpes Zoster:* 1000 mg of **SHILOVA TABLETS** to be taken three times per day for seven days.

*Recurrent genital herpes:* The recommended dosage for the treatment of recurrent genital herpes is 500 mg twice daily for 5 days. Dosing should begin as early as possible. For recurrent episodes of herpes simplex, this should ideally be during the prodromal period or immediately the first signs or symptoms appear. There are no data on the effectiveness of **SHILOVA TABLETS** when initiated more than 24 hours after the onset of signs and symptoms.

*For the prevention (suppression) of recurrences of herpes simplex infection:*

*Immunocompetent patients:* 500 mg to be taken once daily. Some patients with very frequent recurrences (e.g. 10 or more per year) may gain additional benefit from the daily dose of 500 mg being taken as a divided dose (250 mg twice daily).

*Immunocompromised patients:* 500 mg twice daily.

*Prophylaxis of cytomegalovirus infection (CMV) and disease:*

*Adults and adolescents (from 12 years of age):* 2000 mg to be taken four times a day. Dosing should be initiated as early as possible post-transplant. This dose should be reduced according to creatinine clearance (See **Dosage in renal impairment**). The duration of treatment will usually be 90 days but may need to be extended in high-risk patients.

**Paediatric population**

No data are available.

**Special populations**

**Dosage in the elderly:**

Dosage modification is not required unless renal function is impaired (see **Dosage in renal impairment**). Adequate hydration should be maintained.

**Dosage in renal impairment:**

The dose of **SHILOVA TABLETS** should be modified as follows in patients with significantly impaired renal function:

<b>HERPES ZOSTER Creatinine Clearance</b>	<b>SHILOVA TABLETS Dose</b>
15 - 30 ml/min	1000 mg twice a day
< 15 ml/min	1000 mg once a day

<b>RECURRENT GENITAL HERPES Creatinine Clearance</b>	<b>SHILOVA TABLETS Dose</b>
> 15 ml/min	500 mg twice daily
0 - 15 ml/min	500 mg once daily

PREVENTION OF RECURRENCE Creatinine Clearance	SHILOVA TABLETS Dose	
	Immunocompetent	Immunocompromised
15 - 30 ml/min	No dosage adjustment required	No dosage adjustment required
< 15 ml/min	250 mg once a day	500 mg once daily

In patients on haemodialysis, the **SHILOVA TABLETS** dose recommended for patients with a creatinine clearance of less than 15 ml/min should be used, but the dose should be administered after the haemodialysis has been performed.

**CMV PROPHYLAXIS:** The dosage of **SHILOVA TABLETS** should be adjusted in patients with impaired renal function as shown in the table below:

Creatinine Clearance	SHILOVA TABLETS Dose
≥ 75 ml/min	2000 mg four times daily
50 to < 75 ml/min	1500 mg four times daily
25 to < 50 ml/min	1500 mg three times daily
10 to < 25 ml/min	1500 mg twice daily
< 10 ml/min or dialysis**	1500 mg once daily

\*\* In patients on haemodialysis, the **SHILOVA TABLETS** dosage should be administered after the haemodialysis has been performed.

The creatinine clearance should be monitored frequently, especially during periods when renal function is changing rapidly e.g. immediately after transplantation or engraftment. The **SHILOVA TABLETS** dosage should be adjusted accordingly.

**Dosage in hepatic impairment:**

Dose modification is not required in patients with mild or moderate cirrhosis (hepatic synthetic function maintained). Pharmacokinetic data in patients with advanced cirrhosis (impaired hepatic synthetic function and evidence of portal-systemic shuttling) do not indicate the need for dosage

adjustment; however, clinical experience is limited. For higher doses recommended for CMV prophylaxis see section 4.8.

#### **Method of administration**

To be taken orally.

#### **4.3 Contraindications**

**SHILOVA TABLETS** is contraindicated in patients known to be hypersensitive to valaciclovir, aciclovir or any component of their formulations.

Safety in pregnancy and lactation has not been established. (See section 4.6 “).

#### **4.4 Special warnings and precautions for use**

Thrombotic, thrombocytopenic purpura/haemolytic uraemic syndrome (TTP / HUS):

TTP / HUS in some cases resulted in death, has been reported in patients with advanced HIV disease and also in bone marrow transplant and renal transplant recipients.

Cases of acute renal failure have been reported in:

- Elderly patients with or without reduced renal function. Caution should be exercised when administering **SHILOVA TABLETS** to elderly patients, and dosage reduction is recommended for those with impaired renal function.
- Patients with underlying renal disease who received higher than recommended doses of **SHILOVA TABLETS** for their level of renal function. Dosage reduction is recommended when administering **SHILOVA TABLETS** to patients with renal impairment.
- Patients receiving other nephrotoxic medicines. Caution should be exercised when administering **SHILOVA TABLETS** to patients receiving potentially nephrotoxic medicines.
- Patients without adequate hydration. Precipitation of aciclovir in renal tubules may occur when solubility (2,5 mg/ml) is exceeded in the intratubular fluid. Adequate hydration should be maintained for all patients.

In the event of acute renal failure and anuria, the patient may benefit from haemodialysis until renal function is restored.

### **Central Nervous System (CNS)**

CNS adverse reactions, including agitation, hallucinations, confusion, delirium, seizures, and encephalopathy, have been reported in both adult and paediatric patients with or without reduced renal function and in patients with underlying renal disease who received higher than recommended doses of **SHILOVA TABLETS** for their level of renal function. Elderly patients are more likely to have CNS adverse reactions. **SHILOVA TABLETS** should be discontinued if CNS adverse reactions occur.

### **Skin and appendage**

Severe cutaneous adverse reactions (SCARS) including toxic epidermal necrolysis (TEN) Stevens Johnson Syndrome (SJS) and drug reaction with no eosinophilia and systemic symptoms (DRESS), which could be life-threatening or fatal, have been reported with **SHILOVA TABLETS** (see section 4.8). At the time of prescription, patients should be advised of the signs and symptoms of severe skin reactions and closely monitored. If signs and symptoms suggestive of these reactions appear, **SHILOVA TABLETS** should be discontinued immediately, and an alternative treatment should be considered. If the patient has developed a serious reaction such as SJS, TEN or DRESS with use of **SHILOVA TABLETS**, treatment with **SHILOVA TABLETS** must not be restarted in this patient at any time.

### **Use in genital herpes**

Patients should be advised to avoid intercourse when symptoms are present even if treatment with an antiviral has been initiated. Continuous therapy with **SHILOVA TABLETS** in patients with recurrent genital herpes reduces the risk of transmitting genital herpes. It does not cure genital herpes or completely eliminate the risk of transmission. In addition to therapy with valaciclovir, it is recommended that patients use safer sex practices. As genital herpes is a sexually transmitted infection, patients should, in order to further reduce the risk of infecting partners, avoid contact with lesions, damaged skin/mucosa, and avoid intercourse when lesions and/or symptoms are present. Genital herpes is frequently transmitted in the absence of symptoms through asymptomatic viral shedding; therefore, patients should be counselled to use safer sex practices. The effect of **SHILOVA**

**TABLETS** on transmission of sexually transmitted infections other than herpes (including HIV, gonorrhoea, syphilis and Chlamydia) is unknown.

The efficacy of **SHILOVA TABLETS** for reducing transmission of genital herpes has not been established in individuals with multiple partners, non-heterosexual couples and couples not counselled to use safer sex practises.

#### **4.5 Interaction with other medicines and other forms of interaction**

No clinically significant interactions have been identified.

Cimetidine and probenecid increase the area under the plasma concentration time curve of aciclovir by reducing its renal clearance: however, no dosage adjustment is necessary because of the wide therapeutic index of aciclovir. Other medicines which affect renal physiology could affect plasma levels of aciclovir.

In patients receiving high-dose **SHILOVA TABLETS** (8 g/day) for CMV prophylaxis, caution is required during concurrent administration with medicines which compete with aciclovir for elimination, because of the potential for increased plasma levels of one or both medicines or their metabolites.

Increases in plasma AUCs of aciclovir and of the inactive metabolite of mycophenolate mofetil, an immunosuppressant agent used in transplant patients, have been shown when the medicines are co-administered.

Care is also required (with monitoring for changes in renal function) if administering high-dose **SHILOVA TABLETS** with medicines which affect other aspects of renal physiology (e.g. cyclosporin, tacrolimus).

#### **4.6 Fertility, pregnancy and lactation**

Safety in pregnancy and lactation has not been established (see section 4.3).

##### **Teratogenicity:**

Foetal abnormalities were observed in rats.

#### 4.7 Effects on ability to drive and use machines

When driving vehicles or using machines it should be taken into account that dizziness or weariness or visual disturbances may occur.

#### 4.8 Undesirable effects

MedDRA system organ class	Frequency	Adverse reactions
Blood and the lymphatic system disorders	Less frequent	Aplastic anaemia Thrombocytopenia Haemolytic anaemia, micro-angiopathic, neutropenia. Leukocytoclastic vasculitis.
Immune system disorders	Less frequent	Acute hypersensitivity reactions and anaphylaxis and angioedema
Nervous system disorders	Frequent:	Headache
	Less frequent	Decreased consciousness in patients with renal insufficiency Dizziness, somnolence, convulsions Fatigue Aggressive behaviour Agitation Ataxia, coma, confusion, dysarthria, encephalopathy, seizures and tremors
Psychiatric disorders:	Less frequent	Mania and psychosis, including auditory and visual hallucinations.
Eye disorders:	Less frequent	Visual abnormalities
Cardiac disorders:	Less frequent	Tachycardia
Vascular disorder:	Less frequent	Hypertension
Gastrointestinal disorders:	Frequent	Nausea.

**Applicant/PHCR:**

**Product proprietary name:**

**Dosage form and strength:**

**AUROGEN SOUTH AFRICA (PTY) LTD**

**SHILOVA 500 mg/ 1000 mg TABLETS**

**TABLETS 500 mg/ 1000 mg**

**Date: 13/02/2025**

	Less frequent	Gastrointestinal disturbances such as constipation, diarrhoea, loss of appetite, stomach pain, vomiting.
Hepato-biliary disorders:	Less frequent	Hepatitis with liver function test abnormalities, jaundice.
Skin and subcutaneous tissue disorders:	Less frequent	Facial oedema. Skin reactions such as erythema multiforme, photosensitivity, or rash (redness of the skin). Alopecia, Stevens-Johnson syndrome, toxic epidermal necrolysis, urticaria, pruritus.
	Unknown frequency	Acute generalised exanthematous pustulosis (AGEP), Drug reaction with eosinophilia and systemic symptoms (DRESS).
Musculoskeletal, connective tissue and bone disorders:	Less frequent	Arthralgia.
Renal and urinary disorders:	Less frequent	Renal insufficiency manifested by increased serum creatinine, acute renal failure
Reproductive system and breast disorders:	Less frequent	Dysmenorrhoea

### **Reporting of suspected adverse reactions**

Reporting suspected adverse reactions after authorisation of the medicine is important. It allows continued monitoring of the benefit/risk balance of the medicine. Health care providers are requested to report any suspected adverse drug reactions to SAHPRA via the Med Safety APP (Medsafety X SAHPRA) and eReporting platform (who-umc.org) found on SAHPRA website.

### **4.9 Overdose**

#### **Symptoms and signs:**

There are at present no data available on overdose with **SHILOVA TABLETS**.

In animal studies, large doses caused obstructive uropathy and crystalluria.

#### **Management:**

In the event of asymptomatic **SHILOVA TABLETS** overdose occurring, aciclovir is removable by haemodialysis.

## **5 PHARMACOLOGICAL PROPERTIES**

### **5.1 Pharmacodynamic properties**

Pharmacological classification: A.20.2.8 Antiviral agents

Pharmacotherapeutic group: Antivirals for systemic use. Nucleosides and nucleotides excluding reverse transcriptase inhibitors.

ATC Code: J05AB11

Valaciclovir, an antiviral, is the L-valine ester of aciclovir. Aciclovir is a purine (guanine) nucleoside analogue.

Valaciclovir is a nucleoside analogue DNA polymerase inhibitor.

Valaciclovir is rapidly and almost completely converted in man to aciclovir and L-valine probably by the enzyme valaciclovir hydrolase.

Aciclovir is a specific inhibitor of the herpes viruses with *in vitro* activity against herpes simplex viruses (HSV) type 1 and type 2, varicella zoster virus (VZV), Epstein-Barr virus (EBV),

cytomegalovirus (CMV) and human herpes virus 6 (HHV-6). Aciclovir inhibits herpes virus DNA synthesis once it has been phosphorylated to the active triphosphate form. The first stage of phosphorylation requires the activity of a virus-specific enzyme. In the case of HSV, VZV and EBV this enzyme is the viral thymidine kinase (TK), which is only present in virus infected cells.

Selectivity is maintained in CMV with phosphorylation, at least in part, being mediated through the phosphotransferase gene product of the UL97 gene of CMV. This gene encodes for the viral kinase which facilitates the intracellular anabolism of aciclovir.

The requirement for activation of aciclovir by a virus-specific enzyme largely explains its unique selectivity. The phosphorylation process is completed (conversion from mono- to triphosphate) by cellular kinases.

Aciclovir triphosphate competitively inhibits the virus DNA polymerase and incorporation of this nucleoside analogue result in obligate chain termination, halting virus DNA synthesis and thus blocking virus replication.

Extensive monitoring of clinical isolates from patients receiving aciclovir therapy or prophylaxis has revealed that herpes simplex virus and varicella zoster virus with reduced sensitivity to aciclovir is rare in the immunocompetent and is only found infrequently in severely immunocompromised individuals e.g. solid organ or bone marrow transplant recipients, patients receiving chemotherapy for malignant disease and people infected with the human immunodeficiency virus (HIV).

Resistance is normally due to a thymidine kinase deficient phenotype which results in a virus which is profoundly disadvantaged in the natural host. Infrequently, reduced sensitivity to aciclovir has been described as a result of subtle alterations in either the virus thymidine kinase or DNA polymerase. The virulence of these variants resembles that of the wild-type virus.

## **5.2 Pharmacokinetic properties**

### **Absorption:**

After oral administration valaciclovir is well absorbed from the gastrointestinal tract and almost completely converted to aciclovir and L-valine by first-pass intestinal and / or hepatic metabolism.

### **Metabolism:**

Aciclovir is converted to a small extent to inactive metabolites by aldehyde oxidase and by alcohol and aldehyde dehydrogenase. Neither valaciclovir nor aciclovir is metabolised by cytochrome P450 enzymes. Peak plasma concentrations of unconverted valaciclovir are low and transient, generally becoming non-quantifiable 3 hours after administration.

Peak plasma concentration of valaciclovir occurs at median time of 45 – 60 minutes post dose. Peak plasma valaciclovir concentrations are generally less than 0,5 µg/ ml at all doses. After single-dose administration of 1 gram of valaciclovir, average plasma valaciclovir concentrations observed were 0,5; 0,4 and 0,8 µg/ml in patients with hepatic dysfunction, renal insufficiency, and in healthy volunteers who received concomitant cimetidine and probenecid, respectively.

#### **Distribution:**

The binding of valaciclovir to human plasma proteins ranges from 13, 5 % to 17, 9 %. The binding of aciclovir to human plasma proteins ranges from 9 % to 33 %.

#### **Elimination:**

The elimination plasma half-life of aciclovir after both single and multiple dosing with valaciclovir averaged 2, 5 to 3, 3 hours in volunteers with normal renal function.

Less than 1 % of the administered dose of valaciclovir is recovered in the urine. Valaciclovir is eliminated principally as aciclovir and the known aciclovir metabolite, 9-carboxymethoxymethyl-guanine (CMMG), in the urine.

Renal impairment: Following administration of valaciclovir to patients with renal impairment, the average half-life of aciclovir is approximately 14 hours. During haemodialysis, the aciclovir half-life is approximately 4 hours. Approximately one third of aciclovir in the body is removed by dialysis during a 4-hour haemodialysis session. Apparent plasma clearance of aciclovir in dialysis patients is 86, 3 +/- 21, 3 ml/min/1, 73 m<sup>2</sup> compared with 679, 16 +/- 162, 76 ml /min/1, 73 m<sup>2</sup> in healthy volunteers.

#### **Characteristics in patients:**

Herpes zoster and herpes simplex does not significantly alter the pharmacokinetics of valaciclovir and aciclovir after oral administration of **SHILOVA TABLETS**. In transplant recipients receiving

**Applicant/PHCR:** AUROGEN SOUTH AFRICA (PTY) LTD  
**Product proprietary name:** SHILOVA 500 mg/ 1000 mg TABLETS  
**Dosage form and strength:** TABLETS 500 mg/ 1000 mg

**Date:** 13/02/2025

valaciclovir 2000 mg 4 times daily, aciclovir peak concentrations are similar to or greater than those in healthy volunteers receiving the same dose. The estimated daily area under the plasma concentration curves (AUCs) is appreciably greater.

### 5.3 Preclinical safety data

Not applicable

## 6 PHARMACEUTICAL PARTICULARS

### 6.1 List of excipients

- Cellulose microcrystalline,
- Crospovidone, magnesium stearate and
- Povidone.

Opadry blue 13b50578 film coating ingredients:

- Fd&c blue #2/ indigo carmine aluminium lake (c.i. no.: 73015),
- Hypromellose,
- Macrogol,
- Polysorbate 80 and
- Titanium dioxide (c.i. no.: 77891).

### 6.2 Incompatibilities

Not applicable.

### 6.3 Shelf life

36 months

**Applicant/PHCR:** AUROGEN SOUTH AFRICA (PTY) LTD  
**Product proprietary name:** SHILOVA 500 mg/ 1000 mg TABLETS  
**Dosage form and strength:** TABLETS 500 mg/ 1000 mg

**Date:** 13/02/2025

#### 6.4 Special precautions for storage

Store at or below 25 °C.

Keep HDPE containers tightly closed.

Keep the blisters in the carton until required for use.

#### 6.5 Nature and contents of container SHILOVA 500 mg TABLETS:

##### 1) Blister Pack:

Tablets are packed in clear 250 µm PVC film coated with 90 g/m<sup>2</sup> PVdc and printed 25 µm aluminium foil with 7g/m<sup>2</sup> heat seal lacquer.

##### Pack sizes:

**10s** - Each blister contains 5 tablets. Each carton contains 2 blisters of 5 tablets each.

**30s** - Each blister contains 5 tablets. Each carton contains 6 blisters of 5 tablets each.

**42s** – Each blister contains 7 tablets. Each carton contains 6 blisters of 7 tablets each.

##### 2) HDPE Container Pack:

Tablets are packed in 60 ml HDPE white opaque colour container of 38 mm neck finish with 38 mm – 400 RS closure with induction sealing wad. Each container contains 42 tablets.

**Pack size: 42s** - One HDPE container contains 42 tablets.

#### SHILOVA 1 g TABLETS:

##### 1) Blister Pack:

Tablets are packed in Clear 250 µm PVC film coated with 90 g/m<sup>2</sup> PVdC and Printed 25 µm aluminium foil with 7 g/m<sup>2</sup> heat seal lacquer.

**Pack size: 21s** – Each carton contains 3 blisters of 7 tablets each.

##### 2) HDPE Container Pack:

Tablets are packed in 60 ml white opaque colour HDPE container of 38 mm neck finish with 38 mm – 400 RS closure with induction sealing wad. Each container contains 21 tablets.

**Pack size: 21s** - One HDPE container contains 21 tablets.

**Applicant/PHCR:** AUROGEN SOUTH AFRICA (PTY) LTD  
**Product proprietary name:** SHILOVA 500 mg/ 1000 mg TABLETS  
**Dosage form and strength:** TABLETS 500 mg/ 1000 mg

**Date:** 13/02/2025

## **7 HOLDER OF CERTIFICATE OF REGISTRATION**

Aurogen South Africa (Pty) Ltd

Woodhill Office Park, Building 1, First floor,

53 Phillip Engelbrecht Avenue,

Meyersdal, Ext. 12,

Johannesburg,

South Africa.

+27 11 867 9100

[www.auro-space.com](http://www.auro-space.com)

## **8 REGISTRATION NUMBER(S)**

**SHILOVA 500 mg TABLETS:** 45/20.2.8/0590

**SHILOVA 1 g TABLETS:** 45/20.2.8/0591

## **9 DATE OF FIRST AUTHORISATION/ RENEWAL OF THE AUTHORISATION**

15 August 2013

## **DATE OF FIRST AUTHORISATION/ RENEWAL OF THE AUTHORISATION**

13 February 2025