

## **PROFESSIONAL INFORMATION**

### **SCHEDULING STATUS**

Schedule 1

### **PROPRIETARY NAME (AND DOSAGE FORM)**

STREPSILS NUMBING (LOZENGE)

### **COMPOSITION**

Each Lozenge contains

2,4 Dichlorobenzyl Alcohol	1,2 mg
Amylmetacresol	0,6 mg
Lignocaine HCL	10,0 mg

### **PHARMACOLOGICAL ACTION**

STREPSILS NUMBING lozenges have antibacterial and local anaesthetic properties.

### **INDICATIONS**

Symptomatic relief of a sore throat.

### **CONTRA-INDICATIONS**

Hypersensitivity to any of the ingredients.

### **DOSAGE AND DIRECTIONS FOR USE**

Dissolve one lozenge slowly in the mouth every 2 to 3 hours. Not more than 8 lozenges to be sucked in any 24 hours. Not recommended for children under 12 years.

Lignocaine may impair swallowing, and increase the risk of aspiration. Local anaesthetics should be used with caution patients with epilepsy, impaired cardiac conduction, shock or with liver damage, patients with myasthenia gravis are particularly susceptible to the effects of local anaesthetics.

When a sore throat persists for more than two days, or is accompanied by headache, nausea or vomiting, consult your doctor.

If pregnant or breast feeding consult your doctor before using this product. Do not exceed the stated dose.

### **KNOWN SYMPTOMS OF OVERDOSAGE AND PARTICULARS OF ITS TREATMENT**

Treatment is supportive and symptomatic.

### **IDENTIFICATION**

A pale blue-green circular lozenge with a characteristic flavour.

### **PRESENTATION**

Cartons containing 16 blister packed lozenges.

### **STORAGE INSTRUCTIONS**

KEEP OUT OF REACH OF CHILDREN.

Store below 25°C

### **REGISTRATION NUMBER**

28/16.4/0647

### **NAME AND BUSINESS ADDRESS OF THE HOLDER OF CERTIFICATE OF REGISTRATION**

Reckitt Benckiser Pharmaceuticals (Pty) Ltd

8 Jet Park Road Elandsfontein

1601

### **DATE OF PUBLICATION OF THIS PACKAGE INSERT**

24 February 1995