

## **SCHEDULING STATUS**

**S4**

### **1. NAME OF THE MEDICINE**

**XADAGO 50 mg film-coated tablets**

**XADAGO 100 mg film-coated tablets**

### **2. QUALITATIVE AND QUANTITATIVE COMPOSITION**

**XADAGO 50 mg**

Each film-coated tablet contains safinamide methanesulfonate equivalent to 50 mg safinamide.

**XADAGO 100 mg**

Each film-coated tablet contains safinamide methanesulfonate equivalent to 100 mg safinamide.

Sugar free.

For the full list of excipients, see section 6.1.

### **3. PHARMACEUTICAL FORM**

Film-coated tablets

**XADAGO 50 mg**

Orange to copper, round, biconcave, film-coated tablet of 7 mm diameter with metallic gloss, embossed with the strength "50" on one side of the tablet.

**XADAGO 100 mg**

Orange to copper, round, biconcave, film-coated tablet of 9 mm diameter with metallic gloss, embossed with the strength "100" on one side of the tablet.

### **4. CLINICAL PARTICULARS**

#### **4.1 Therapeutic indications**

XADAGO is indicated for the treatment of adult patients with idiopathic Parkinson's disease (PD) as add-on therapy to a stable dose of levodopa (L-dopa) alone or in combination with other PD medicines in mid-to late-stage fluctuating patients.

#### **4.2 Posology and method of administration**

## PROFESSIONAL INFORMATION

---

### **Posology**

Treatment with XADAGO should be started at 50 mg per day. This daily dose may be increased to 100 mg/day on the basis of individual clinical need.

If a dose is missed the next dose should be taken at the usual time the next day.

### *Elderly patients*

No change in dose is required for elderly patients.

Experience of use of safinamide in patients over 75 years of age is limited.

### *Hepatic impairment*

XADAGO use in patients with severe hepatic impairment is contraindicated (see section 4.3). No dose adjustment is required in patients with mild hepatic impairment. The lower dose of 50 mg/day is recommended for patients with moderate hepatic impairment. If patients progress from moderate to severe hepatic impairment, XADAGO should be stopped (see section 4.4).

### *Renal impairment*

No change in dose is required for patients with renal impairment.

### *Paediatric population*

The safety and efficacy of XADAGO in children and adolescents under 18 years of age have not been established. No data are available.

### **Method of administration**

For oral use.

XADAGO should be taken with water.

XADAGO may be taken with or without food.

### **4.3 Contraindications**

- Hypersensitivity to safinamide or to any of the excipients (see section 6.1).
- Concomitant treatment with other monoamine oxidase (MAO) inhibitors (see sections 4.4 and 4.5).
- Concomitant treatment with pethidine (see sections 4.4 and 4.5).
- Use in patients with severe hepatic impairment (see section 4.2).
- Use in patients with albinism, retinal degeneration, uveitis, inherited retinopathy or severe

progressive diabetic retinopathy (see sections 4.4 and 5.3).

#### **4.4 Special warnings and precautions for use**

##### **General warning**

In general, safinamide may be used with selective serotonin re-uptake inhibitors (SSRIs) at the lowest effective dose, with caution for serotonergic symptoms. In particular, the concomitant use of safinamide and fluoxetine or fluvoxamine should be avoided, or if concomitant treatment is necessary these medicines should be used at low doses (see section 4.5). A washout period corresponding to 5 half-lives of the SSRI used previously should be considered prior to initiating treatment with safinamide.

At least 7 days must elapse between discontinuation of safinamide and initiation of treatment with MAO inhibitors or pethidine (see section 4.3 and 4.5).

When safinamide is co-administered with products that are BCRP substrates, please refer to the professional information (PI) for that particular medicine.

##### **Hepatic impairment**

Caution should be exercised when initiating treatment with safinamide in patients with moderate hepatic impairment. In case patients progress from moderate to severe hepatic impairment, treatment with XADAGO should be stopped (see sections 4.2, 4.3 and 5.2).

##### **Potential for retinal degeneration in patients with prior history of retinal disease**

XADAGO should not be administered to patients with ophthalmological history that would put them at increased risk for potential retinal effects (e.g., family history of hereditary retinal disease, or history of uveitis) see sections 4.3 and 5.3.

##### **Impulse control disorders (ICDs)**

Impulse control disorders can occur in patients treated with dopamine agonists and/or dopaminergic treatments. Some reports of ICDs have also been observed with other MAO-inhibitors. Safinamide treatment has not been associated with any increase in the appearance of ICDs.

Patients and carers should be made aware of the behavioural symptoms of ICDs that were observed in patients treated with MAO-inhibitors, including cases of compulsions, obsessive thoughts, pathological gambling, increased libido, hypersexuality, impulsive behaviour and

compulsive spending or buying.

### **Dopaminergic side effects**

Safinamide used as an adjunct to levodopa may potentiate the side effects of levodopa, and pre-existing dyskinesia may be exacerbated, requiring a decrease of levodopa (refer to levodopa Professional Information for further information). This effect was not seen when safinamide was used as an adjunct to dopamine agonists in early-stage PD patients.

In addition, Dopamine dysregulation syndrome (DDS) has been reported in literature reports in patients treated with carbidopa/levodopa. Although Safinamide (contained in XADAGO) never showed this risk both alone nor when administered with carbidopa/levodopa, patients and caregivers should be warned of the potential risk of developing DDS when Xadago is co-administered with carbidopa/levodopa.

Dopamine Dysregulation Syndrome (DDS) is an addictive disorder seen in some patients treated with carbidopa/levodopa. Affected patients show a compulsive pattern of dopaminergic drug misuse above doses adequate to control motor symptoms, which may in some cases result in severe dyskinesias.

## **4.5 Interaction with other medicines and other forms of interaction**

### ***In vivo and in vitro pharmacodynamic interactions***

#### *MAO inhibitors and pethidine*

XADAGO must not be administered along with other MAO inhibitors (including moclobemide) as there may be a risk of non-selective MAO inhibition that may lead to a hypertensive crisis (see section 4.3).

Serious adverse reactions have been reported with the concomitant use of pethidine and MAO inhibitors. As this may be a class-effect, the concomitant administration of safinamide and pethidine is contraindicated (see section 4.3).

There have been reports of interactions with the concomitant use of MAO inhibitors and sympathomimetic medicines. In view of the MAO inhibitory activity of safinamide, concomitant administration of safinamide and sympathomimetics, such as those present in nasal and oral decongestants or cold and flu medicines containing ephedrine or pseudoephedrine, requires caution (see section 4.4).

#### *Dextromethorphan*

## PROFESSIONAL INFORMATION

---

There have been reports of interactions with the concomitant use of dextromethorphan and non-selective MAO inhibitors. In view of the MAO inhibitory activity of safinamide, the concomitant administration of safinamide and dextromethorphan is not recommended, or if concomitant treatment is necessary, it should be used with caution (see section 4.4).

### *Antidepressants*

The concomitant use of XADAGO and fluoxetine or fluvoxamine should be avoided (see section 4.4), this precaution is based on the occurrence of serious adverse reactions (e.g. serotonin syndrome), although rare, that have occurred when SSRIs and dextromethorphan have been used with MAO inhibitors (including selective MAO-B inhibitors), selective serotonin reuptake inhibitors (SSRIs), serotonin-norepinephrine reuptake inhibitors (SNRIs), tricyclic antidepressants, tetracyclic antidepressants, triazolopyridine antidepressants, cyclobenzaprine, opioid drugs and methylphenidate, amphetamine and their derivatives. If necessary, the concomitant use of these medicines should be at the lowest effective dose. A washout period corresponding to 5 half-lives of the SSRI used previously should be considered prior to initiating treatment with XADAGO. Serotonin syndrome symptoms may include mental status changes (e.g., agitation, hallucinations, delirium, and coma), autonomic instability (e.g., tachycardia, labile blood pressure, dizziness, diaphoresis, flushing, and hyperthermia), neuromuscular symptoms (e.g., tremor, rigidity, myoclonus, hyperreflexia, and incoordination), seizures, and/or gastrointestinal symptoms (e.g., nausea, vomiting, and diarrhoea).

Serious adverse reactions have been reported with the concomitant use of selective serotonin reuptake inhibitors (SSRIs), serotonin norepinephrine reuptake inhibitors (SNRIs), tricyclic/tetracyclic antidepressants and MAO inhibitors (see section 4.4). In view of the selective and reversible MAO-B inhibitory activity of safinamide, antidepressants may be administered but used at the lowest doses necessary.

### ***In vivo and in vitro pharmacokinetic interactions***

Safinamide may transiently inhibit BCRP *in vitro*. In interaction studies in human, a weak interaction was observed with rosuvastatin (AUC increase between 1,25 and 2,00-fold) but no significant interaction was found with diclofenac. It is recommended to monitor patients when safinamide is taken with medicinal products that are BCRP substrates (e.g., rosuvastatin, pitavastatin, pravastatin, ciprofloxacin, methotrexate, topotecan, diclofenac or glyburide) and to refer to their professional information to determine if a dose adjustment is needed.

Safinamide is almost exclusively eliminated via metabolism, largely by high capacity amidases

## PROFESSIONAL INFORMATION

---

that have not yet been characterised. Sildenafil is eliminated mainly in the urine. In human liver microsomes (HLM), the N-dealkylation step appears to be catalysed by CYP3A4, as sildenafil clearance in HLM was inhibited by ketoconazole by 90 %.

Sildenafil inhibits OCT1 *in vitro* at clinically relevant portal vein concentrations. Therefore, caution is necessary when sildenafil is taken concomitantly with medicines that are OCT1 substrates and have a  $t_{max}$  similar to sildenafil (2 hours) (e.g. metformin, aciclovir, ganciclovir) as exposure to these substrates might be increased as a consequence.

The metabolite NW-1153 is a substrate for OAT3 at clinically relevant concentrations. Medicines that are inhibitors of OAT3 given concomitantly with XADAGO may reduce clearance of NW-1153, i.e., and thus may increase its systemic exposure. The systemic exposure of NW-1153 is low (1/10 of parent sildenafil). This potential increase is most likely of no clinical relevance as NW-1153, the first product in the metabolic pathway, is further transformed to secondary and tertiary metabolites.

### **Paediatric population**

Interaction studies have only been performed in adults.

## **4.6 Fertility, pregnancy and lactation**

### **Women of childbearing potential**

XADAGO should not be given to women of childbearing potential unless adequate contraception is practiced.

### **Pregnancy**

There are no or limited amount of data from the use of XADAGO in pregnant women. Studies in animals have shown reproductive toxicity (see section 5.3). XADAGO is not recommended during pregnancy and in women of childbearing potential not using contraception.

### **Breastfeeding**

Available pharmacodynamic/toxicological data in animals have shown excretion of sildenafil in milk (for details see 5.3).

A risk for the breastfed child cannot be excluded. XADAGO should not be used during breastfeeding.

### **Fertility:**

## PROFESSIONAL INFORMATION

Animal studies indicate that XADAGO treatment is associated with adverse reactions on female rat reproductive performance and sperm quality. Male rat fertility is not affected (see section 5.3).

### 4.7 Effects on ability to drive and use machines

Somnolence and dizziness may occur during safinamide treatment, therefore patients should be cautioned about using hazardous machines, including motor vehicles, until they are reasonably certain that XADAGO does not affect them adversely.

### 4.8 Undesirable effects

#### Summary of the safety profile

Dyskinesia was the most common adverse reaction reported in XADAGO patients when used in combination with L-dopa alone or in combination with other PD treatments.

Serious adverse reactions are known to occur with the concomitant use of SSRIs, SNRIs, tricyclic/tetracyclic antidepressants and MAO inhibitors, such as hypertensive crisis (high blood pressure, collapse), neuroleptic malignant syndrome (confusion, sweating, muscle rigidity, hyperthermia, CPK increase), serotonin syndrome (confusion, hypertension, muscle stiffness, hallucinations), and hypotension. With MAO-inhibitors there have been reports of interactions with concomitant use of sympathomimetic medicines.

Impulse control disorders: pathological gambling, increased libido, hypersexuality, compulsive spending or buying, binge eating and compulsive eating can occur in patients treated with dopamine agonists and/or other dopaminergic treatments.

#### Tabulated list of adverse reactions

The tabulation below includes all adverse reactions in clinical trials where adverse reactions were considered related.

Adverse reactions are ranked under headings of frequency using the following conventions: very common ( $\geq 1/10$ ), common ( $\geq 1/100$  to  $< 1/10$ ), uncommon ( $\geq 1/1\ 000$  to  $< 1/100$ ), rare ( $\geq 1/10\ 000$  to  $< 1/1\ 000$ ), very rare ( $< 1/10\ 000$ ) and not known (cannot be estimated from the available data).

System Organ Class	Very common	Common	Uncommon	Rare
Infections and infestations			Urinary tract infection	Bronchopneumonia, furuncle, nasopharyngitis,

PROFESSIONAL INFORMATION

				pyoderma, rhinitis, tooth infection, viral infection
Neoplasms benign, malignant and unspecified (incl cysts and polyps)			Basal cell carcinoma	Acrochordon, Melanocytic naevus, Seborrheic keratosis, skin papilloma
Blood and lymphatic system disorders			Anaemia, leukopenia, red blood cell abnormality	Eosinophilia, lymphopenia
Metabolism and nutrition disorders			Decreased appetite, hypertriglyceridaemia, increased appetite, hypercholesterolaemia, hyperglycaemia	Cachexia, hyperkalaemia
Psychiatric disorders		Insomnia	Hallucination, depression, abnormal dreams, anxiety, confusional state, affect lability, libido increased, psychotic disorder,	Compulsions, delirium, disorientation, illusion, impulsive behaviour, loss of libido, obsessive thoughts,

PROFESSIONAL INFORMATION

			restlessness, sleep disorder	paranoia, premature ejaculation, sleep attacks, social phobia, suicidal ideation
Nervous system disorders		Dyskinesia somnolence, dizziness, headache, Parkinson's disease	Paraesthesia, balance disorder, hypoesthesia, dystonia, head discomfort, dysarthria, syncope, cognitive disorder	Coordination abnormal, disturbance in attention, dysgeusia, hyporeflexia, radicular pain, Restless Legs Syndrome, sedation
Eye disorders		Cataract	Vision blurred, scotoma, diplopia, photophobia, retinal disorder, conjunctivitis, glaucoma	Amblyopia, chromatopsia, diabetic retinopathy, erythroptosis, eye haemorrhage, eye pain, eyelid oedema, hypermetropia, keratitis, lacrimation increased, night blindness, papilloedema, presbyopia, strabismus

## PROFESSIONAL INFORMATION

Ear and labyrinth disorders			Vertigo	
Cardiac disorders			Palpitations, tachycardia, sinus bradycardia, dysrhythmia	Myocardial infarction
Vascular disorders		Orthostatic hypotension	Hypertension, hypotension, varicose vein	Arterial spasm, arteriosclerosis, hypertensive crisis
Respiratory, thoracic and mediastinal disorders			Cough, dyspnoea, rhinorrhoea	Bronchospasm, dysphonia, oropharyngeal pain, oropharyngeal spasm
Gastrointestinal disorders		Nausea	Constipation, dyspepsia, vomiting, dry mouth, diarrhoea, abdominal pain, gastritis, flatulence, abdominal distension, salivary hypersecretion, gastroesophageal reflux disease, aphthous stomatitis	Peptic ulcer, retching, upper gastrointestinal haemorrhage

Date of approval: 24 January 2023

PROFESSIONAL INFORMATION

Hepatobiliary disorders				Hyperbilirubinaemia
Skin and subcutaneous tissue disorders			Hyperhidrosis, pruritus generalised, photosensitivity reaction, erythema	Alopecia, blister, dermatitis contact, dermatosis, ecchymosis, lichenoid keratosis, night sweats, pain of skin, pigmentation disorder, psoriasis, seborrhoeic dermatitis
Musculoskeletal and connective tissue disorders			Back pain, arthralgia, muscle spasms, muscle rigidity, pain in extremity, muscular weakness, sensation of heaviness	Ankylosing spondylitis, flank pain, joint swelling, musculoskeletal pain, myalgia, neck pain, osteoarthritis, synovial cyst
Renal and urinary disorders			Nocturia, dysuria	Micturition urgency, polyuria, pyuria, urinary hesitation
Reproductive system and breast disorders			Erectile dysfunction	Benign prostatic hyperplasia, breast disorder, breast pain

Date of approval: 24 January 2023

PROFESSIONAL INFORMATION

<p>General disorders and administration site conditions</p>			<p>Fatigue, asthenia, gait disturbance, oedema peripheral, pain, feeling hot</p>	<p>Drug effect decreased, drug intolerance, feeling cold, malaise, pyrexia, xerosis</p>
<p>Investigations</p>			<p>Weight decreased, weight increased, blood creatine phosphokinase increased, blood triglycerides increased, blood glucose increased, blood urea increased, blood alkaline phosphatase increased, blood bicarbonate increased, blood creatinine increased, electrocardiogram QT prolonged, liver function test abnormal, urine analysis abnormal, blood pressure increased, blood</p>	<p>Blood calcium decreased, blood potassium decreased, blood cholesterol decreased, body temperature increased, cardiac murmur, cardiac stress test abnormal, haematocrit decreased, haemoglobin decreased, international normalised ratio decreased, lymphocyte count decreased, platelet count decreased, very low density lipoprotein increased</p>

## PROFESSIONAL INFORMATION

			pressure decreased, ophthalmic diagnostic procedures abnormal	
Injury, poisoning and procedural complications		Fall	Foot fracture	Contusion, fat embolism, head injury, mouth injury, skeletal injury
Social circumstances				Gambling

### **Description of selected adverse reactions**

Dyskinesia occurred early in treatment, was rated “severe”, led to discontinuation in very few patients (approx. 1,5 %), and did not require reduction of dose in any patient.

### **Post marketing experience**

The following adverse reactions have been identified during post approval use of safinamide. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to estimate their frequency or establish a causal relationship to drug exposure.

Immune system disorders: Hypersensitivity: A post marketing report describes a patient who developed a hypersensitivity reaction consisting of swelling of the tongue and gingiva, dyspnea and skin rash. The symptoms resolved shortly after XADAGO was discontinued, but reappeared following rechallenge a month later.

### **Reporting of suspected adverse reactions:**

Reporting suspected adverse reactions after authorisation of XADAGO is important. It allows continued monitoring of the benefit/risk balance of XADAGO. Health care providers are asked to report any suspected adverse reactions via the “**6.04 Adverse Drug Reaction Reporting Form**”, found online under SAHPRA’s publications: <https://www.sahpra.org.za/Publications/Index/8>.

#### **4.9 Overdose**

In one patient suspected of consuming more than the daily prescribed dose of 100 mg for one month, symptoms of confusion, sleepiness, forgetfulness and dilated pupils were reported. These symptoms resolved on discontinuing XADAGO, without sequelae.

The expected pattern of events or symptoms following intentional or accidental overdose with safinamide would be those related to its pharmacodynamic profile: MAO-B inhibition with activity-dependent inhibition of Na<sup>+</sup> channels. The symptoms of an excessive MAO-B inhibition (increase in dopamine level) could include hypertension, postural hypotension, hallucinations, agitation, nausea, vomiting, and dyskinesia.

There is no known antidote to XADAGO or any specific treatment for XADAGO overdose. If an important overdose occurs, XADAGO treatment should be discontinued and supportive treatment should be administered as clinically indicated.

### **5. PHARMACOLOGICAL PROPERTIES**

#### **5.1 Pharmacodynamic properties**

Category and class: A 5.4.1 Anti-Parkinsonism preparations.

Pharmacotherapeutic group: Anti-Parkinson-drugs, monoamine oxidase-B inhibitors.

ATC code: N04BD03

#### **Mechanism of action**

Safinamide acts through both dopaminergic and non-dopaminergic mechanisms of action.

Safinamide is a highly selective and reversible MAO-B inhibitor causing an increase in extracellular levels of dopamine in the striatum. Safinamide is associated with state-dependent inhibition of voltage-gated sodium (Na<sup>+</sup>) channels, and modulation of stimulated release of glutamate. To what extent the non-dopaminergic effects contribute to the overall effect has not been established.

#### **Pharmacodynamic effects**

Population PK models developed from studies in patients with Parkinson's disease indicate that the pharmacokinetic and pharmacodynamics effects of safinamide were not dependent on age, gender, weight, renal function and exposure to levodopa, indicating that dose adjustments will not be required based on these variables.

## PROFESSIONAL INFORMATION

---

Pooled analyses of adverse event data from placebo-controlled studies in Parkinson's disease patients indicate that the concomitant administration of safinamide together with a broad category of commonly used medicines in this patient population (antihypertensive, beta-blockers, cholesterol lowering, non-steroidal anti-inflammatory medicines, proton pump inhibitors, antidepressants, etc.) was not associated with an increased risk for adverse events. Studies were not stratified for co-medication, and no randomised interaction studies were performed for these medicines.

### **Clinical efficacy**

#### *Studies in mid- to late-stage PD patients*

The efficacy of safinamide as add-on treatment in mid-to late-stage PD (LSPD) patients with motor fluctuations, currently receiving L-dopa alone or in combination with other PD medicines, was evaluated in two double-blind, placebo-controlled studies: Study SETTLE (Study 27919; 50 – 100 mg/day; 24 weeks), and Study 016/018 (50 and 100 mg/day; 2-year, double-blind, placebo-controlled study).

The primary efficacy parameter was the change from baseline to endpoint in ON Time without troublesome dyskinesia.

Secondary efficacy parameters included OFF Time, UPDRS II and III (Unified Parkinson's Disease Rating Scale – sections II and III), and CGI-C (Clinical Global Impression of Change).

Both the SETTLE and 016/018 studies indicated significant superiority of safinamide, compared to placebo, at the target doses of 50 and 100 mg/day for the primary, and selected secondary, efficacy variables, as summarized in the table below. The effect on ON Time was maintained at the end of the 24-month double-blind treatment period for both safinamide doses as compared to placebo.

PROFESSIONAL INFORMATION

Study	016 (24 weeks)			016/018 (2 years)			27919 (SETTLE) (24 weeks)	
	Placebo	Safinamide		Placebo	Safinamide		Placebo	Safinamide
		50	100		50	100		
<b>Randomized</b>	222	223	224	222	223	224	275	274
<b>Age (years) (b)</b>	59,4 (9,5)	60,1 (9,7)	60,1 (9,2)	59,4 (9,5)	60,1 (9,7)	60,1 (9,2)	62,1 (9,0)	61,7 (9,0)
<b>PD Duration (years) (b)</b>	8,4 (3,8)	7,9 (3,9)	8,2 (3,8)	8,4 (3,8)	7,9 (3,9)	8,2 (3,8)	9,0 (4,9)	8,9 (4,4)
<b>ON time without troublesome dyskinesia (hrs) (c)</b>								
Baseline (b)	9,3 (2,2)	9,4 (2,2)	9,6 (2,5)	9,3 (2,2)	9,4 (2,2)	9,6 (2,5)	9,1 (2,5)	9,3 (2,4)
Change LSM (SE)	0,5 (0,2)	1,0 (0,2)	1,2 (0,2)	0,8 (0,2)	1,4 (0,2)	1,5 (0,2)	0,6 (0,1)	1,4 (0,1)
LS Diff vs Placebo		0,5	0,7		0,6	0,7		0,9
95% CI		[0,1, 0,9]	[0,3, 1,0]		[0,1, 1,0]	[0,2, 1,1]		[0,6, 1,2]
p-value		0,0054	0,0002		0,011 0	0,0028		<0,000 1
<b>OFF time (hrs) (c)</b>								
Baseline (b)	5,3 (2,1)	5,2 (2,0)	5,2 (2,2)	5,3 (2,1)	5,2 (2,2)	5,2 (2,1)	5,4 (2,0)	5,3 (2,0)
Change LSM (SE)	-0,8 (0,20)	-1,4 (0,20)	-1,5 (0,20)	-1,0 (0,20)	-1,5 (0,19)	-1,6 (0,19)	-0,5 (0,10)	-1,5 (0,10)
LS Diff vs Placebo		-0,6	-0,7		-0,5	-0,6		-1,0
95% CI		[-0,9, -0,3]	[-1,0, -0,4]		[-0,8, -0,2]	[-0,9, -0,3]		[-1,3, -0,7]

PROFESSIONAL INFORMATION

		-0,3]	-0,4]		-0,2]	-0,3]		-0,7]
p-value		0,0002	<0,000 1		0,002 8	0,0003		<0,000 1
<b>UPDRS III (c)</b>								
Baseline (b)	28,6 (12,0)	27,3 (12,8)	28,4 (13,5)	28,6 (12,0)	27,3 (12,8)	28,4 (13,5)	23,0 (12,8)	22,3 (11,8)
Change LSM (SE)	-4,5 (0,83)	-6,1 (0,82)	-6,8 (0,82)	-4,4 (0,85)	-5,6 (0,84)	-6,5 (0,84)	-2,6 (0,34)	-3,5 (0,34)
LS Diff vs Placebo		-1,6	-2,3		-1,2	-2,1		-0,9
95% CI		[-3,0, -0,2]	[-3,7, -0,9]		[-2,6, 0,2]	[-3,5, -0,6]		[-1,8, 0,0]
p-value		0,0207	0,0010		0,093 9	0,0047		0,0514
<b>UPDRS II (c)</b>								
Baseline (b)	12,2 (5,9)	11,8 (5,7)	12,1 (5,9)	12,2 (5,9)	11,8 (5,7)	12,1 (5,9)	10,4 (6,3)	10,0 (5,6)
Change LSM (SE)	-1,2 (0,4)	-1,9 (0,4)	-2,3 (0,4)	-1,4 (0,3)	-2,0 (0,3)	-2,5 (0,3)	-0,8 (0,2)	-1,2 (0,2)
LS Diff vs Placebo		-0,7	-1,1		-0,6	-1,1		-0,4
95% CI		[-1,3, -0,0]	[-1,7, -0,5]		[-1,3, 0,0]	[-1,8, -0,4]		[-0,9, 0,0]
p-value		0,0367	0,0007		0,067 6	0,0010		0,0564
<b>Responder analyses (post-hoc) (e) n(%)</b>								
ON time increase ≥60 minutes	93 (43,9)	119 (54,8)	121 (56,0)	100 (47,2)	125 (57,6)	117 (54,2)	116 (42,5)	152 (56,3)
p-value		0,0233	0,0122		0,030 8	0,1481		0,0013

## PROFESSIONAL INFORMATION

≥60 minutes increase ON time and decrease in OFF time and ≥ 30% improvement UPDRS III	32 (15,1)	52 (24,0)	56 (25,9)	28 (13,2)	43 (19,8)	42 (19,4)	24 (8,8)	49 (18,1)
p-value		0,0216	0,0061		0,067 1	0,0827		0,0017
CGI-C: patients who were much/very much improved	42 (19,8)	72 (33,2)	78 (36,1)	46 (21,7)	62 (28,6)	64 (29,6)	26 (9,5)	66 (24,4)
p-value (f)		0,0017	0,0002		0,096 2	0,0575		<0,000 1
<p>(a) Daily targeted dose, (b) Mean (SD), (c) analysis population (mITT); MMRM model for change from Baseline to Endpoint includes treatment, region, and visit as fixed effects, and baseline value as a covariate; (d) target dose of 100 mg/day; (e) analysis population (mITT); data are presented as the number (percentage) of patients in each group meeting the responder definition (f) chi-square test of the odds ratio of the treatment groups compared to placebo using a logistic regression model, with fixed effects for treatment and country.</p> <p>SE Standard Error, SD Standard deviation, LSM Least Square Mean, LS Diff. Least Square Difference vs Placebo</p> <p>mITT Population: Study 016/018 - Placebo (n=212), safinamide 50 mg/day (n=217) and 100 mg/day (n=216), and SETTLE - Placebo (n=270), safinamide 50-100 mg/day (n=273).</p>								

### Paediatric population

The pharmacodynamic effects of safinamide have not been assessed in children and adolescents.

## 5.2 Pharmacokinetic properties

### Absorption

Safinamide absorption is rapid after single and multiple oral dosing, reaching  $T_{max}$  in the time range 1,8 – 2,8 h after dosing under fasting conditions. Absolute bioavailability is high (95 %), showing that safinamide is almost completely absorbed after oral administration and first pass

## PROFESSIONAL INFORMATION

---

metabolism is negligible. The high absorption classifies safinamide as a highly permeable substance.

A slight delay in  $T_{max}$  was observed in the fed state relative to the fasted condition, but there was no effect on safinamide  $AUC_{0-\infty}$  or  $C_{max}$ . XADAGO may be administered with or without food.

### **Distribution**

The volume of distribution ( $V_{ss}$ ) is approximately 165 L which is 2,5-fold of body volume indicating extensive extravascular distribution of safinamide. Total clearance was determined to be 4,6 L/h classifying safinamide as a low clearance substance.

Plasma protein binding of safinamide is 88 – 90 %.

### **Biotransformation**

In humans, safinamide is almost exclusively eliminated via metabolism (urinary excretion of unchanged safinamide was < 10 %) mediated principally through high-capacity amidases, that have not yet been characterized. *In vitro* experiments indicated that inhibition of amidases in human hepatocytes led to complete suppression of the NW-1153 formation. Amidase present in blood, plasma, serum, simulated gastric fluid and simulated intestinal fluid as well as human carboxylesterases hCE-1 and hCE-2 are not responsible for the biotransformation of safinamide to NW-1153. The amidase FAAH was able to catalyse the formation of NW-1153 at low rates only. Therefore, other amidases are likely to be involved in the conversion to NW-1153. Safinamide's metabolism is not dependent on cytochrome P450 (CYP) based enzymes.

Metabolite structure elucidation revealed three metabolic pathways of safinamide. The principal pathway involves hydrolytic oxidation of the amide moiety leading to the primary metabolite safinamide acid (NW-1153). Another pathway involves oxidative cleavage of the ether bond forming *O*-debenzylated safinamide (NW-1199). Finally, the *N*-dealkylated acid (NW-1689) is formed by oxidative cleavage of the amine bond of either safinamide (minor) or the primary safinamide acid metabolite (NW-1153) (major). The *N*-dealkylated acid (NW-1689) undergoes conjugation with glucuronic acid yielding its acyl glucuronide. None of these metabolites are pharmacologically active.

Safinamide does not appear to significantly induce or inhibit enzymes at clinically relevant systemic concentrations. *In vitro* metabolism studies have indicated that there is no meaningful

## PROFESSIONAL INFORMATION

---

induction or inhibition of cytochrome P450, CYP2A6, 2B6, 2C8, 2C9, 2C19, 2D6, 2E1 and 3A3/5 at concentrations which are relevant ( $C_{max}$  of free safinamide 0,4  $\mu\text{M}$  at 100 mg/day) in man. Dedicated interaction studies performed with ketoconazole, L-dopa and CYP1A2 and CYP3A4 substrates (caffeine and midazolam), did not detect any clinically significant effects on the pharmacokinetics of safinamide, or L-dopa, caffeine and midazolam.

A mass balance study showed that the plasma  $\text{AUC}_{0-24\text{h}}$  of the unchanged  $^{14}\text{C}$ -safinamide accounted for approximately 30 % of the total radioactivity  $\text{AUC}_{0-24\text{h}}$ , indicative of an extensive metabolism.

### Transporters

Preliminary *in vitro* studies have shown that safinamide is not a substrate for the transporters P-gp, BCRP, OAT1B1, OAT1B3, OATP1A2 or OAT2P1. Metabolite NW-1153 is not a substrate for OCT2, or OAT1, but it is substrate for OAT3. This interaction has the potential to reduce the clearance of NW-1153 and increase its exposure; however the systemic exposure of NW-1153 is low (1/10 of parent safinamide), and as it is metabolised to secondary and tertiary metabolites, it is unlikely to be of any clinical relevance.

Safinamide transiently inhibits BCRP in the small intestine (see section 4.5). At concentrations of 50  $\mu\text{M}$ , safinamide inhibited OATP1A2 and OATP2P1. The relevant plasma concentrations of safinamide are substantially lower, therefore a clinically relevant interaction with co-administered substrates of these transporters is unlikely. NW-1153 is not an inhibitor of OCT2, MATE1, or MATE2-K up to concentrations of 5  $\mu\text{M}$ .

### Linearity/non-linearity

The pharmacokinetics of safinamide are linear after single and repeated doses. No time-dependency was observed.

### Elimination

Safinamide undergoes almost complete metabolic transformation (< 10 % of the administered dose was found unchanged in urine). Substance-related radioactivity was largely excreted in urine (76 %) and only to a low extent in faeces (1,5 %) after 192 hours. The terminal elimination half-life of total radioactivity was approximately 80 hours.

## PROFESSIONAL INFORMATION

---

The elimination half-life of safinamide is 20 – 30 hours. Steady state is reached within one week.

### **Hepatic impairment**

Safinamide exposure in patients with mild hepatic disease increased marginally (30 % in AUC), while in patients with moderate hepatic impairment exposure increased by approximately 80 % (see section 4.2).

### **Renal impairment:**

Moderate or severe renal impairment did not alter the exposure to safinamide, compared to healthy subjects (see section 4.2).

### **5.3 Preclinical safety data**

Retinal degeneration and loss of photoreceptor cells were observed in albino and pigmented rats after repeated safinamide oral dosing resulting in systemic exposure below the anticipated systemic exposure in patients given the maximal therapeutic dose. In albino rats administered safinamide orally for two years, retinal scarring and cataracts were observed at all doses tested. No retinal degeneration was noted in monkeys despite higher systemic exposure than in rodents or in patients at the maximum human dose.

Long-term studies in animals have shown convulsions (1,6 to 12,8 times human clinical exposure, based on plasma AUC). Liver hypertrophy and fatty changes were seen only in rodent livers at exposures similar to humans. Phospholipidosis was seen mainly in the lungs in rodents (at exposures similar to humans) and monkeys (at exposures greater than 12-fold higher than human).

#### *Genotoxicity*

Safinamide did not present genotoxic potential in *in vivo* and in several *in vitro* systems using bacteria or mammalian cells.

#### *Carcinogenicity*

The results obtained from carcinogenicity studies in mice and rats showed no evidence of tumorigenic potential related to safinamide at systemic exposures up to 2,3 to 4,0 times respectively, the anticipated systemic exposure in patients given the maximal therapeutic dose.

## PROFESSIONAL INFORMATION

---

### *Impairment of fertility*

Fertility studies in female rats showed reduced number of implantations and corpora lutea at exposures in excess of 3 times the anticipated human exposure. Male rats showed minor abnormal morphology and reduced speed of sperm cells at exposures in excess of 1,4 times the anticipated human exposure. Male rat fertility was not affected.

### *Embryofoetal Studies*

In embryo-foetal developmental studies in rats and rabbits, malformations were induced at safinamide exposures 2 and 3-fold above human clinical exposure, respectively. The combination of safinamide with levodopa/carbidopa (80/20 mg/kg/day) resulted in additive effects in the embryo-foetal development studies with a higher incidence of foetal skeletal abnormalities than seen with either treatment alone.

In a pre- and postnatal developmental rat study, pup mortality, absence of milk in the stomach and neonatal hepatotoxicity were observed at dose levels similar to the anticipated clinical exposure. Toxic effects on the liver and accompanying symptoms as yellow/orange skin and skull, in pups exposed to safinamide during lactation are mediated mainly via in utero exposure, whereas exposure via the mother's milk had only a minor influence.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

#### **Tablet core**

Crospovidone type A

Magnesium stearate

Microcrystalline cellulose

Silica, colloidal anhydrous

#### **Tablet coating**

Hypromellose

Iron oxide red (E172)

Macrogol (6000)

Mica (E555)

Titanium dioxide (E171)

**6.2 Incompatibilities**

Not applicable.

**6.3 Shelf life**

3 years.

**6.4 Special precautions for storage**

Store at or below 30 °C.

Keep blister strips in carton until required for use.

**6.5 Nature and contents of container**

PVC/PVDC/Aluminium blister packs of 14, 28, 30, 90 or 100 tablets.

Not all pack sizes may be marketed.

**6.6 Special precautions for disposal and other handling**

No special requirements for disposal.

**7. HOLDER OF CERTIFICATE OF REGISTRATION**

Adcock Ingram Limited

1 New Road

Erand Gardens

Midrand 1685,

South Africa

Customer Care: 0860 ADCOCK / 232625

**8. REGISTRATION NUMBERS**

XADAGO 50 mg: 55/5.4.1/0806

XADAGO 100 mg: 55/5.4.1/0807

**9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

24 January 2023

**10. DATE OF REVISION OF THE TEXT**

## PROFESSIONAL INFORMATION

---

Date of approval: 24 January 2023